

ISSN 2224-5227

2016 • 1

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
БАЯНДАМАЛАРЫ

ДОКЛАДЫ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН

REPORTS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN

ЖУРНАЛ 1944 ЖЫЛДАН ШЫҒА БАСТАҒАН
ЖУРНАЛ ИЗДАЕТСЯ С 1944 г.
PUBLISHED SINCE 1944



Бас редактор
ҚР ҰҒА академигі **М.Ж. Жұрынов**

Редакция алқасы:

хим.ғ. докторы, проф., ҚР ҰҒА академигі **Әдекенов С.М.** (бас редактордың орынбасары), эк.ғ. докторы, проф., ҚР ҰҒА академигі **Әділов Ж.М.**, мед. ғ. докторы, проф., ҚР ҰҒА академигі **Арзықұлов Ж.А.**, техн. ғ. докторы, проф., ҚР ҰҒА академигі **Бишімбаев У.К.**, а.-ш.ғ. докторы, проф., ҚР ҰҒА академигі **Есполов Т.И.**, техн. ғ. докторы, проф., ҚР ҰҒА академигі **Мұтанов Г.М.**, физ.-мат.ғ. докторы, проф., ҚР ҰҒА академигі **Өтелбаев М.О.**, пед. ғ. докторы, проф., ҚР ҰҒА академигі **Пралиев С.Ж.**, геогр.ғ. докторы, проф., ҚР ҰҒА академигі **Северский И.В.**; тарих.ғ. докторы, проф., ҚР ҰҒА академигі **Сыдықов Е.Б.**, физ.-мат.ғ. докторы, проф., ҚР ҰҒА академигі **Тәкібаев Н.Ж.**, физ.-мат.ғ. докторы, проф., ҚР ҰҒА академигі **Харин С.Н.**, тарих ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Әбүсейітова М.Х.**, экон. ғ. докторы, проф., ҰҒА корр. мүшесі **Бейсембетов И.К.**, биол. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Жамбакин К.Ж.**, тарих ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Кәрібаев Б.Б.**, мед. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Локшин В.Н.**, геол.-мин. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Өмірсеріков М.Ш.**, физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Рамазанов Т.С.**, физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Садыбеков М.А.**, хим.ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Сатаев М.И.**; ҚР ҰҒА құрметті мүшесі, а.-ш.ғ. докторы, проф. **Омбаев А.М.**

Редакция кеңесі:

Украинаның ҰҒА академигі **Гончарук В.В.** (Украина), Украинаның ҰҒА академигі **Неклюдов И.М.** (Украина), Беларусь Республикасының ҰҒА академигі **Гордиенко А.И.** (Беларусь), Молдова Республикасының ҰҒА академигі **Дука Г.** (Молдова), Тәжікстан Республикасының ҰҒА академигі **Илолов М.И.** (Тәжікстан), Қырғыз Республикасының ҰҒА академигі **Эркебаев А.Э.** (Қырғызстан), Ресей ҒА корр. мүшесі **Величкин В.И.** (Ресей Федерациясы); хим.ғ. докторы, профессор **Марек Сикорски** (Польша), тех.ғ. докторы, профессор **Потапов В.А.** (Украина), биол.ғ. докторы, профессор **Харун Парлар** (Германия), профессор **Гао Энджун** (КХР), филос. ғ. докторы, профессор **Стефано Перни** (Ұлыбритания), ғ. докторы, профессор **Богуслава Леска** (Польша), философия ғ. докторы, профессор **Полина Прокопович** (Ұлыбритания), профессор **Вуйцик Вольдемар** (Польша), профессор **Нур Изура Удзир** (Малайзия), д.х.н., профессор **Нараев В.Н.** (Ресей Федерациясы)

Главный редактор
академик НАН РК **М.Ж. Журинов**

Редакционная коллегия:

доктор хим. наук, проф., академик НАН РК **С.М. Адекенов** (заместитель главного редактора), доктор экон. наук, проф., академик НАН РК **Ж.М. Адилов**, доктор мед. наук, проф., академик НАН РК **Ж.А. Арзыкулов**, доктор техн. наук, проф., академик НАН РК **В.К. Бишимбаев**, доктор сельскохоз. наук, проф., академик НАН РК **Т.И. Есполов**, доктор техн. наук, проф., академик НАН РК **Г.М. Мутанов**, доктор физ.-мат. наук, проф., академик НАН РК **М.О. Отелбаев**, доктор пед. наук, проф., академик НАН РК **С.Ж. Пралиев**, доктор геогр. наук, проф., академик НАН РК **И.В. Северский**, доктор ист. наук, проф., академик НАН РК **Е.Б. Сыдыков**, доктор физ.-мат. наук, проф., академик НАН РК **Н.Ж. Такибаев**, доктор физ.-мат. наук, проф., академик НАН РК **С.Н. Харин**, доктор ист. наук, проф., чл.-корр. НАН РК **М.Х. Абусейтова**, доктор экон. наук, проф., чл.-корр. НАН РК **И.К. Бейсембетов**, доктор биол. наук, проф., чл.-корр. НАН РК **К.Ж. Жамбакин**, доктор ист. наук, проф., чл.-корр. НАН РК **Б.Б. Карибаев**, доктор мед. наук, проф., чл.-корр. НАН РК **В.Н. Локшин**, доктор геол.-мин. наук, проф., чл.-корр. НАН РК **М.Ш. Омирсериков**, доктор физ.-мат. наук, проф., чл.-корр. НАН РК **Т.С. Рамазанов**, доктор физ.-мат. наук, проф., чл.-корр. НАН РК **М.А. Садыбеков**, доктор хим. наук, проф., чл.-корр. НАН РК **М.И. Сатаев**; почетный член НАН РК, доктор сельскохоз. наук, проф., **А.М. Омбаев**

Редакционный совет:

академик НАН Украины **Гончарук В.В.** (Украина), академик НАН Украины **И.М. Неклюдов** (Украина), академик НАН Республики Беларусь **А.И. Гордиенко** (Беларусь), академик НАН Республики Молдова **Г. Дука** (Молдова), академик НАН Республики Таджикистан **М.И. Илолов** (Таджикистан), член-корреспондент РАН **Величкин В.И.** (Россия); академик НАН Кыргызской Республики **А.Э. Эркебаев** (Кыргызстан), д.х.н., профессор **Марек Сикорски** (Польша), д.т.н., профессор **В.А. Потапов** (Украина), д.б.н., профессор **Харун Парлар** (Германия), профессор **Гао Энджун** (КНР), доктор философии, профессор **Стефано Перни** (Великобритания), доктор наук, профессор **Богуслава Леска** (Польша), доктор философии, профессор **Полина Прокопович** (Великобритания), профессор **Вуйчик Вольдемар** (Польша), профессор **Нур Изура Удзир** (Малайзия), д.х.н., профессор **В.Н. Нараев** (Россия)

«Доклады Национальной академии наук Республики Казахстан» ISSN 2224-5227

Собственник: Республиканское общественное объединение «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №5540-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год. Тираж: 2000 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 218-220, тел. 272-13-19, 272-13-18

<http://nauka-nanrk.kz>, reports-science.kz

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

©Национальная академия наук Республики Казахстан, 2016 г.

Editor-in-chief

M.Zh. Zhurinov, academician of NAS RK

Editorial board:

S.M. Adekenov (deputy editor in chief), Doctor of Chemistry, prof., academician of NAS RK; **Zh.M. Adilov**, Doctor of Economics, prof., academician of NAS RK; **Zh.A. Arzykulov**, Doctor of Medicine, prof., academician of NAS RK; **V.K. Bishimbayev**, Doctor of Engineering, prof., academician of NAS RK; **T.I. Yespolov**, Doctor of Agriculture, prof., academician of NAS RK; **G.M. Mutanov**, Doctor of Physics and Mathematics, prof., academician of NAS RK; **M.O. Otelbayev**, Doctor of Physics and Mathematics, prof., academician of NAS RK; **S.Zh. Praliyev**, Doctor of Education, prof., academician of NAS RK; **I.V. Seversky**, Doctor of Geography, prof., academician of NAS RK; **Ye.B. Sydykov**, Doctor of Historical Sciences, prof., academician of NAS RK; **N.Zh. Takibayev**, Doctor of Physics and Mathematics, prof., academician of NAS RK; **S.N. Kharin**, Doctor of Physics and Mathematics, prof., academician of NAS RK; **M.Kh. Abuseitova**, Doctor of Historical Sciences, prof., corr. member of NAS RK; **I.K. Beisembetov**, Doctor of Economics, prof., corr. member of NAS RK; **K.Zh. Zhambakin**, Doctor of Biological Sciences, prof., corr. member of NAS RK; **B.B. Karibayev**, Doctor of Historical Sciences, prof., corr. member of NAS RK; **V.N. Lokshin**, Doctor of Medicine, prof., corr. member of NAS RK; **M.Sh. Omirserikov**, Doctor of Geology and Mineralogy, prof., corr. member of NAS RK; **T.S. Ramazanov**, Doctor of Physics and Mathematics, prof., corr. member of NAS RK; **M.A. Sadybekov**, Doctor of Physics and Mathematics, prof., corr. member of NAS RK; **M.I. Satayev**, Doctor of Chemistry, prof., corr. member of NAS RK; **A.M. Ombayev**, Honorary Member of NAS RK, Doctor of Agriculture, prof.

Editorial staff:

V.V. Goncharuk, NAS Ukraine academician (Ukraine); **I.M. Neklyudov**, NAS Ukraine academician (Ukraine); **A.I. Gordienko**, NAS RB academician (Belarus); **G. Duca**, NAS Moldova academician (Moldova); **M.I. Iolov**, NAS Tajikistan academician (Tajikistan); **A.E. Erkebayev**, NAS Kyrgyzstan academician (Kyrgyzstan); **V.I. Velichkin**, RAS corr.member (Russia); **Marek Sikorski**, Doctor of Chemistry, prof. (Poland); **V.A. Potapov**, Doctor of Engineering, prof. (Ukraine); **Harun Parlar**, Doctor of Biological Sciences, prof. (Germany); **Gao Endzhun**, prof. (PRC); **Stefano Perni**, Doctor of Philosophy, prof. (UK); **Boguslava Leska**, dr, prof. (Poland); **Pauline Prokopovich**, Doctor of Philosophy, prof. (UK); **Wójcik Waldemar**, prof. (Poland), **Nur Izura Udzir**, prof. (Malaysia), **V.N. Narayev**, Doctor of Chemistry, prof. (Russia)

Reports of the National Academy of Sciences of the Republic of Kazakhstan.

ISSN 2224-5227

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 5540-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 2000 copies

Editorial address: 28, Shevchenko str., of.219-220, Almaty, 050010, tel. 272-13-19, 272-13-18,

<http://nauka-nanrk.kz/> reports-science.kz

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

© National Academy of Sciences of the Republic of Kazakhstan, 2016

D. Suragan

Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan.
E-mail: suragan@list.ru

A NONLOCAL BOUNDARY VALUE PROBLEM ON THE HEISENBERG GROUP

Abstract. In this note we construct a well-posed boundary value problem for the Kohn Laplacian with nonlocal boundary conditions in a bounded smooth domain on the Heisenberg group. It is also showed that the boundary value problem is explicitly solvable for any bounded smooth domain.

Keywords: nonlocal boundary value problem, Kohn Laplacian, Heisenberg group.

Introduction. In a bounded domain of the Euclidean space $\Omega \in \mathbb{R}^d, d \geq 2$, it is very well known that the solution to the Laplacian equation

$$\Delta u(x) = f(x), \quad x \in \Omega, \quad (1)$$

is given by the Green formula (or the Newton potential formula)

$$u(x) = \int_{\Omega} \varepsilon_d(x-y) f(y) dy, \quad x \in \Omega, \quad (2)$$

for suitable functions f supported in Ω . Here ε_d is the fundamental solution to Δ in \mathbb{R}^d given by

$$\varepsilon_d(x-y) = \begin{cases} \frac{1}{(2-d)\omega_d} |x-y|^{d-2}, & d \geq 3, \\ \frac{1}{2\pi} \log|x-y|, & d = 2, \end{cases} \quad (3)$$

where $\omega_d = \frac{2\pi^{d/2}}{\Gamma(\frac{d}{2})}$ is the surface area of the unit sphere in \mathbb{R}^d .

An interesting question having several important applications is what boundary conditions can be put on u on the (smooth) boundary $\partial\Omega$ so that equation (1) complemented by this boundary condition would have the solution in Ω still given by the same formula (2), with the same kernel ε_d given by (3). It turns out that the answer to this question is the integral boundary condition

$$-\frac{1}{2}u(x) + \int_{\partial\Omega} \frac{\partial \varepsilon_d(x-y)}{\partial n_y} u(y) dS_y - \int_{\partial\Omega} \varepsilon_d(x-y) \frac{\partial u(y)}{\partial n_y} dS_y = 0, \quad x \in \partial\Omega, \quad (4)$$

where $\frac{\partial}{\partial n_y}$ denotes the outer normal derivative at a point y on $\partial\Omega$. A converse question to the one above would be to determine the trace of the Newton potential (2) on the boundary surface $\partial\Omega$, and one can use the potential theory to show that it has to be given by (4).

The boundary condition (4) appeared in M. Kac's work [10] where he called it and the subsequent spectral analysis the principle of not feeling the boundary". This was further expanded in Kac's book [11] with several further applications to the spectral theory and the asymptotics of the Weyl's eigenvalue counting function. In [12] by using the boundary condition (1.4) the eigenvalues and eigenfunctions of the

Newton potential (2) were explicitly calculated in the 2-disk and in the 3-ball. In general, the boundary value problem (1)-(4) has various interesting properties and applications (see, for example, Kac [10, 11] and Saito [19]). The boundary value problem (1)-(4) can also be generalized for higher degrees of the Laplacian, see [13, 14].

In this note we are interested in and give answers to the following questions:

– What happens if an elliptic operator (the Laplacian) in (1) is replaced by a hypoelliptic operator? We will realize this as a model of replacing the Euclidean space by the Heisenberg group and the Laplacian on \mathbb{R}^d by a sub-Laplacian (or the Kohn-Laplacian) on \mathbb{H}_{n-1} . We will show that the boundary condition (4) is replaced by the integral boundary condition (15) in this setting (see also (11)).

– Since the theory of boundary value problems for elliptic operators is well understood, we know that the single condition (4) on the boundary $\partial\Omega$ of a bounded domain Ω guarantees the unique solvability of the equation (1) in Ω . Is this uniqueness preserved in the hypoelliptic model as well for a suitably chosen replacement of the boundary condition (4)? The case of the second order operators is favourable from this point of view due to the validity of the maximum principle, see Bony [1]. The Dirichlet problem has been considered by Jerison [9]. The answer in the case of the boundary value problem in our setting is given in Theorem 2.1.

We now describe the setting of this paper. The Heisenberg group \mathbb{H}_{n-1} is the space $\mathbb{C}^{n-1} \times \mathbb{R}$ with the group operation given by

$$(\zeta, t) \circ (\eta, \tau) = (\zeta + \eta, t + \tau + 2\operatorname{Im}\zeta\eta), \quad (5)$$

for $(\zeta, t), (\eta, \tau) \in \mathbb{C}^{n-1} \times \mathbb{R}$. Writing $\zeta = x + iy$, with $x_j, y_j, j = 1, \dots, n-1$, the real coordinates on \mathbb{H}_{n-1} , the left-invariant vector fields

$$\begin{aligned} \tilde{X}_j &= \frac{\partial}{\partial x_j} + 2y_j \frac{\partial}{\partial t}, j = 1, \dots, n-1, \\ \tilde{Y}_j &= \frac{\partial}{\partial y_j} - 2x_j \frac{\partial}{\partial t}, j = 1, \dots, n-1, \\ T &= \frac{\partial}{\partial t} \end{aligned}$$

form a basis for the Lie algebra \mathfrak{h}_{n-1} of \mathbb{H}_{n-1} .

On the other hand, \mathbb{H}_{n-1} can be viewed as the boundary of the Siegel upper half space in \mathbb{C}^n , $\mathbb{H}_{n-1} = \{(\zeta, z_n) \in \mathbb{C}^n : \operatorname{Im} z_n = |\zeta|^2, \zeta = (z_1, \dots, z_{n-1})\}$. Parameterizing \mathbb{H}_{n-1} by $z = (\zeta, t)$, where $\mathbb{H}_{n-1} t = \operatorname{Re} z_n$, a basis for the complex tangent space of \mathbb{H}_{n-1} at the point z is given by the left-invariant vector fields

$$X_j = \frac{\partial}{\partial z_j} + i\bar{z} \frac{\partial}{\partial t}, j = 1, \dots, n-1.$$

We denote their conjugates by $X_{\bar{j}} = X_j = \frac{\partial}{\partial \bar{z}_j} - i z \frac{\partial}{\partial t}$. The operator

$$\square_{a,b} = \sum_{j=1}^{n-1} (aX_j X_{\bar{j}} + bX_{\bar{j}} X_j), a + b = n-1, \quad (6)$$

is a left-invariant, rotation invariant differential operator that is homogeneous of degree two (cf. [3]). This operator is a slight generalisation of the standard sub-Laplacian or Kohn-Laplacian \square_b on the Heisenberg group \mathbb{H}_{n-1} which, when acting on the coefficients of a $(0, q)$ -form can be written as

$$\square_b = -\frac{1}{n-1} \sum_{j=1}^{n-1} ((n-1-q) X_j X_{\bar{j}} + q X_{\bar{j}} X_j).$$

Folland and Stein [4] found that a fundamental solution of the operator $\square_{a,b}$ is a constant multiple of

$$s(z) = s(\xi, t) = \frac{1}{(t+i|\xi|^2)^a (t-i|\xi|^2)^b}, \quad (7)$$

and defined the Newton potential (volume potential) for a function f with compact support contained in a set $\Omega \in \mathbb{H}_{n-1}$ by

$$u(z) = \int_{\Omega} f(\xi) s(t^{-1}z) dv(\xi), \quad (8)$$

with dv being the volume element (the Haar measure on \mathbb{H}_{n-1}), coinciding with the Lebesgue measure on $\mathbb{C}^{n-1} \times \mathbb{R}$. More precisely, they proved that

$$\square_{a,b} u = c_{a,b} f,$$

where the constant $c_{a,b}$ is zero if a and $b = -1, -2, \dots, n, n+1, \dots$, and $c_{a,b} \neq 0$ if a or $b \neq -1, -2, \dots, n, n+1, \dots$. In fact, then we can take

$$c_{a,b} = \frac{2(a^2 + b^2) \text{Vol}(B_1)}{(2i)^n} \frac{(n-1)!}{a(a-1) \dots (a-n+1)} (1 - \exp(-2i\pi))$$

for $a \notin \mathbb{Z}$, see the proof of Theorem 1.6 in Romero [15]. Similar conclusions by a different methods were obtained by Greiner and Stein [8]. For a more general analysis of fundamental solutions for sub-Laplacian we can refer to Folland [5] as well as to a discussion and references in Stein [20].

In the above notation, the distribution $\frac{1}{c_{a,b}} s$ is the fundamental solution of $\square_{a,b}$, while s satisfies the equation

$$\square_{a,b} s = c_{a,b} \delta. \quad (9)$$

However, although we could have rescaled s for it to become the fundamental solution, we prefer to keep the notation yielding (9) in order to follow the notation of [4] and [15] to be able to refer to their results directly.

In this paper we assume that $c_{a,b} \neq 0$, i.e. both a and $b \neq -1, -2, \dots, n, n+1, \dots$. In addition, without loss of generality we may also assume that $a, b \geq 0$.

Now, in analogy to the elliptic boundary value problem (1)-(4) for the Laplacian Δ in \mathbb{R}^d , we consider the hypoelliptic boundary value problem for the sub-Laplacian $\square_{a,b}$ on \mathbb{H}_{n-1} , namely the equation

$$\square_{a,b} u = c_{a,b} f \quad (10)$$

in a bounded set $\Omega \subset \mathbb{H}_{n-1}$ with smooth boundary $\partial\Omega$. The aim of this paper is to find a boundary condition of the Newton potential u on $\partial\Omega$, such that with this boundary condition the equation (10) has a unique solution, which is the Newton potential (8).

Basing our arguments on the analysis of Folland and Stein [4] and Romero [15] we show that the boundary condition (4) for the Laplacian in \mathbb{R}^d is now replaced by the integral boundary condition (5) in this setting, namely by the condition

$$(c_{a,b} - H.R(z))u(z) - \int_{\partial\Omega} s(\xi, z) \langle \nabla^{b,a} u(\xi), dv(\xi) \rangle + p.v.Wu(z) = 0, z \in \partial\Omega,$$

on the boundary $\partial\Omega$, where $H.R(z)$ is the so-called half residue, and where the second and the third term can be interpreted as coming from the suitably defined respectively single and double layer potentials S and W for the problem. See Section 2 for the definitions and the precise statement. In Section 2 by using properties of fundamental solutions we construct a well-posed boundary value problem for the differential equation (10) with the required properties.

The Kohn Laplacian. Let $\Omega \subset \mathbb{H}_{n-1}$ be an open bounded domain with a smooth boundary $\partial\Omega \in C^\infty$. Consider the following analogy of the Newton potential on the Heisenberg group

$$u(z) = \int_{\Omega} f(\xi) \varepsilon(\xi, z) dv(\xi) \quad \text{in } \Omega, \quad (11)$$

where $\varepsilon(\xi, z) = \varepsilon(\xi^{-1}z)$ is the rescaled fundamental solution (1.7) of the sub-Laplacian, satisfying (1.9). As we mentioned u is a solution of (10) in Ω . The aim of this section is to find a boundary condition for u such that with this boundary condition the equation (10) has a unique solution in $C^2(\Omega)$, say, and this solution is the Newton potential (11).

We recall a few notions and properties first. For $z = (\zeta, t) \in \mathbb{H}_{n-1}$, we define its norm by $|z| = (|\zeta|^4 + |t|^2)^{\frac{1}{4}}$. As any (quasi-)norm on \mathbb{H}_{n-1} , this satisfies a triangle inequality with a constant, and allows for a polar decomposition. For $0 < \alpha < 1$, Folland and Stein [4] defined the anisotropic Holder spaces $\Gamma_\alpha\{\Omega\}$ by

$$\Gamma_\alpha\{\Omega\} = \left\{ f: \Omega \rightarrow \mathbb{C} : \sup_{\substack{z_1, z_2 \in \Omega \\ z_1 \neq z_2}} \frac{|f(z_2) - f(z_1)|}{|z_1^{-1}z_2|^\alpha} < \infty \right\}.$$

For $k \in \mathbb{N}$ and $0 < \alpha < 1$, one defines $\Gamma_{k+\alpha}\{\Omega\}$ as the space of all $f: \Omega \rightarrow \mathbb{C}$ such that all complex derivatives of f of order k belong to $\Gamma_\alpha\{\Omega\}$.

A starting point for us will be that if $f \in \Gamma_\alpha(\Omega)$ for $\alpha < 0$ then u defined by (11) is twice differentiable in the complex directions and satisfies the equation $\square_{a,b}u = c_{a,b}f$. We refer to Folland and Stein [4], Greiner and Stein [8], and to Romero [15] for three different approaches to this property. Moreover, Folland and Stein have shown that if $f \in \Gamma_\alpha(\Omega, \text{loc})$ and $\square_{a,b}u = c_{a,b}f$, then $f \in \Gamma_{\alpha+2}(\Omega, \text{loc})$. These results extend those known for the Laplacian, in suitably redefined anisotropic Holder spaces.

We record relevant single and double layer potentials for the problem (10). In [9], Jerison used the single layer potential defined by

$$S_0 g(z) = \int_{\partial\Omega} g(\xi) \varepsilon(\xi, z) dS(\xi),$$

which, however, is not integrable over characteristic points. On the contrary, the functional

$$Sg(z) = \int_{\partial\Omega} g(\xi) \varepsilon(\xi, z) \langle X_j, dv(\xi) \rangle,$$

where $\langle X_j, dv \rangle$ is the canonical pairing between vector fields and differential forms, is integrable over the whole boundary $\partial\Omega$. Moreover, it was shown in [16, Theorem 2.3] that if the density of $g(\xi) \langle X_j, dv \rangle$ in the operator S is bounded then $Sg \in \Gamma_\alpha(\mathbb{H}_{n-1})$ for all $\alpha < 1$. Parallel to S , it is natural to use the operator

$$Wu(z) = \int_{\partial\Omega} u(\xi) \langle \nabla^{b,a} \varepsilon(\xi, z), dv(\xi) \rangle \quad (12)$$

as a double layer potential. Our main result:

Theorem 2.1. Let $\varepsilon(\xi, z) = \varepsilon(\xi^{-1}z)$ be the rescaled fundamental solution to $\square_{a,b}$, so that

$$\square_{a,b} \varepsilon = c_{a,b} \delta \quad \text{on } \mathbb{H}_{n-1} \quad (13)$$

for any $f \in \Gamma_\alpha(\Omega)$, the Newton potential (11) is the unique solution in $C^2(\Omega) \cap C^1(\bar{\Omega})$ of the equation

$$\square_{a,b} u = c_{a,b} f \quad (14)$$

with the boundary condition

$$(c_{a,b} - H.R.(z))u(z) + \lim_{\delta \rightarrow 0} \int_{\partial\Omega \setminus \{|\xi - z| < \delta\}} u(\xi) \nabla^{a,b} \varepsilon(\xi, z), dv(\xi) - \int_{\partial\Omega} \varepsilon(\xi, z), \langle \nabla^{a,b} \varepsilon(\xi, z), dv(\xi) \rangle = 0, \quad \text{для } z \in \partial\Omega, \quad (15)$$

where $H.R.(z)$ is the so-called half residue given by the formula

$$H.R.(z) = \lim_{\delta \rightarrow 0} \int_{\partial\Omega \setminus \{|\xi - z| < \delta\}} \langle \nabla^{a,b} \varepsilon(\xi, z), dv(\xi) \rangle, \quad (16)$$

c

$$\nabla^{a,b} g = \sum_{j=1}^{n-1} (aX_j g X_j + bX_j g X_j).$$

The half residue $H.R.(z)$ in (16) appears in the jump relations for the problem (14) in the following way. The double layer potential Wu in (12) has two limits

$$W^+u(z) = \lim_{z_0 \rightarrow z} \int_{\partial\Omega} u(\xi) \langle \nabla^{a,b} \varepsilon(\xi, z_0), dv(\xi) \rangle$$

and

$$W^-u(z) = \lim_{z_0 \rightarrow z} \int_{\partial\Omega} u(\xi) \langle \nabla^{a,b} \varepsilon(\xi, z_0), dv(\xi) \rangle,$$

and the principal value

$$W^0u(z) = p.v. Wu(z) = \lim_{\delta \rightarrow 0} \int_{\partial\Omega \setminus \{|\xi - z| < \delta\}} u(\xi) \langle \nabla^{a,b} \varepsilon(\xi, z), dv(\xi) \rangle.$$

We note that this principal value enters as the second term in the integral boundary condition (15). It was proved in [16, Theorem 2.4] that for sufficiently regular u (e.g. $u \in \Gamma_\alpha(\Omega)$) and $z \in \partial\Omega$ these limits exist and satisfy the jump relations

$$\begin{aligned} W^+u(z) - W^-u(z) &= c_{a,b} u(z), \\ W^0u(z) - W^-u(z) &= H.R.(z)u(z), \\ W^+u(z) - W^0u(z) &= (c_{a,b} - H.R.(z))u(z), \end{aligned} \quad (17)$$

the last property (17) following from the first two by subtraction.

Proof of Theorem 2.1. Since the solid potential

$$u(z) = \int_{\partial\Omega} u(\xi) \langle \nabla^{b,a} \varepsilon(\xi, z), dv(\xi) \rangle \quad (18)$$

is a solution of (14), from the aforementioned results of Folland and Stein it follows that u is locally in $\Gamma_{\alpha+2}(\Omega, loc)$ and that it is twice complex differentiable in Ω . In particular, it follows that $u \in C^2(\Omega) \cap C^1(\bar{\Omega})$.

The following representation formula can be derived from the generalised second Green's formula (see Theorem 4.5 in [15] and cf. [15]), for $u \in C^2(\Omega) \cap C^1(\bar{\Omega})$ we have

$$\begin{aligned} c_{a,b} u(z) &= c_{a,b} \int_{\Omega} f(\xi) \langle \varepsilon(\xi, z), dv(\xi) \rangle + \int_{\partial\Omega} u(\xi) \langle \nabla^{b,a} \varepsilon(\xi, z), dv(\xi) \rangle \\ &\quad - \int_{\partial\Omega} \varepsilon(\xi) \langle \nabla^{b,a} u(\xi, z), dv(\xi) \rangle \quad \text{for any } z \in \Omega. \end{aligned} \quad (19)$$

Since $u(z)$ given by (18) is a solution of (14), using it in (19) we get

$$\int_{\partial\Omega} u(\xi) \langle \nabla^{a,b} \varepsilon(\xi, z_0), dv(\xi) \rangle - \int_{\partial\Omega} \varepsilon(\xi, z) \langle \nabla^{a,b,a} u(\xi) dv(\xi) \rangle = 0 \quad (20)$$

for any $z \in \Omega$

It is easy to see that the fundamental solution, i.e. the function $s(z)$ in (7) is homogeneous of degree $-2n+2$, that is

$$s(\lambda, z) = \lambda^{-2\alpha-2b} s(z) = \lambda^{-2n+2} s(z) \text{ for any } \lambda > 0,$$

since $\alpha + b = n - 1$. It follows that and its first order complex derivatives are locally integrable. Since $s(\xi, z) = s(\xi^{-1}, z)$, we obtain that as z approaches the boundary, we can pass to the limit in the second term in (20).

By using this and the relation (17) as $z \in \Omega$ approaches the boundary $\partial\Omega$ from inside, we find that

$$(c_{a,b} - H.R(z))u(z) + \lim_{\delta \rightarrow 0} \int_{\partial\Omega \setminus \{|\xi-z| < \delta\}} u(\xi) \langle \nabla^{a,b} s(\xi, z), dv(\xi) \rangle - \int_{\partial\Omega} s(\xi, z) \langle \nabla^{b,a} u(\xi) dv(\xi) \rangle = 0, \text{ for any } z \in \partial\Omega. \quad (21)$$

This shows that (11) is a solution of the boundary value problem (14) with the boundary condition (15).

Now let us prove its uniqueness. If the boundary value problem has two solutions u and u_1 , then the function $w = u - u_1 \in C^2(\Omega) \cap C^1(\overline{\Omega})$ satisfies the homogeneous equation

$$\square_{a,b} w = 0 \text{ in } \Omega, \quad (22)$$

and the boundary condition (15), i.e.

$$c_{a,b} - H.R(z))w(z) + \lim_{\delta \rightarrow 0} \int_{\partial\Omega \setminus \{|\xi-z| < \delta\}} w(\xi) \langle \nabla^{a,b} s(\xi, z), dv(\xi) \rangle - \int_{\partial\Omega} s(\xi, z) \langle \nabla^{b,a} w(\xi) dv(\xi) \rangle = 0 \quad (23)$$

for any $z \in \partial\Omega$.

Since $f \equiv 0$ in this case instead of (19) we have the following representation formula

$$c_{a,b} w(z) = \int_{\partial\Omega} w(\xi, z) \langle \nabla^{a,b} s(\xi) dv(\xi) \rangle - \int_{\partial\Omega} s(\xi, z) \langle \nabla^{b,a} w(\xi) dv(\xi) \rangle \quad (24)$$

for any $z \in \partial\Omega$. As above, by using the properties of the double and single layer potentials as $z \rightarrow \partial\Omega$, we obtain

$$c_{a,b} w(z) = c_{a,b} - H.R(z))w(z) + \lim_{\delta \rightarrow 0} \int_{\partial\Omega \setminus \{|\xi-z| < \delta\}} w(\xi) \langle \nabla^{a,b} s(\xi, z), dv(\xi) \rangle - \int_{\partial\Omega} s(\xi, z) \langle \nabla^{b,a} w(\xi) dv(\xi) \rangle \quad (25)$$

for any $z \in \partial\Omega$. Comparing this with (23) we arrive at

$$w(z) = 0, z \in \partial\Omega. \quad (26)$$

The homogeneous equation (22) with the Dirichlet boundary condition (26) has only trivial solution $w \equiv 0$ in Ω . This shows that the boundary value problem (14) with the boundary condition (15) has a unique solution in $C^2(\Omega) \cap C^1(\overline{\Omega})$. This completes the proof of Theorem 2.1.

REFERENCES

- [1] Bony J.M. Principe du maximum, inegalite de Harnack et unicite du problem de Cauchy pour les operateurs elliptiques degeneres. Ann. Inst. Fourier (Grenoble), 19 (1969), 277-304.
- [2] Fisher V. and Ruzhansky M. Quantization on nilpotent Lie groups, to appear in Progress in Mathematics, Birkhäuser, 2015.
- [3] Folland G.B., Kohn J.J. The Newmann Problem for the Cauchy Riemann Complex, Princeton University Press, Princeton, NJ, 1972.
- [4] Folland G. B. and Stein E. M. Estimates for the $\bar{\partial}_b$ complex and analysis on the Heisenberg group, Comm. Pure Appl. Math. 27 (1974), 429-522.

- [5] Folland G. B. Subelliptic estimates and function spaces on nilpotent Lie groups. Ark. Mat., 13 (1975), 161-207.
- [6] Geller D. Liouville's Theorem for homogeneous groups. Comm. Partial Differential Equations, 8 (1983), 1665-1677.
- [7] Geller D. Analytic pseudodifferential operators for the Heisenberg group and local solvability, mathematical Notes, 37. Princeton University Press, Princeton, NJ, 1990.
- [8] Greiner P.C. and Stein E.M. Estimates for the Neumann problem. Princeton University Press, Princeton, NJ, 1977.
- [9] Jerison D.S. The Dirichlet problem the Kohn Laplacian on the Heisenberg group, I and II. Funct. Anal. 43 (1981), 97-142, 224-257.
- [10] Kac M. on some connections between probability theory and differential and integral equations, Proceedings of the Second Berkeley Symposium on mathematical Statistics and Probability, 1950, university of California Press, Berkeley and Los Angeles, 1951, 189-215.
- [11] Kac M. Integration in function spaces and some of its applications, Lezioni Fermiane. Accademia Nazionale Dei Lincei, Pisa. 1980.
- [12] Kal'menov T.Sh., Suragan D. To spectral problems for the volume potential, Doklady Mathematics, 80 (2009), 646-649.
- [13] Kal'menov T.Sh., Suragan D. Boundary conditions for the volume potential for the polyharmonic equation, differential Equations, 48 (2012), 604-608.
- [14] Kalmenov T. Sh. and Suragan D. A boundary condition and Spectral problems for the Newton Potentials, Operator Theory: Advances and Applications, Vol. 216 (2011), 187-210.
- [15] Romero C. Potential theory for the Kohn Laplacian on the Heisenberg group, PhD thesis, University of Minnesota, 1991.

Д. Сұраған

ҚР БҒМ Математика және математикалық үлгілеу институты, Алматы, Қазақстан

ГЕЙЗЕНБЕРГ ГРУППАСЫНДАҒЫ ЛОКАЛДЫ ЕМЕС ШЕКАРАЛЫҚ ЕСЕП

Аннотация. Жұмыста Гейзенберг группасындағы шеңелген тегіс облыста локалды емес шекаралық шартты Кон Лаплас теңдеуіне қисында шекаралық есебі құрастырылды. Және де кез келген шеңелген тегіс облыста алынған шекаралық есеп айқын түрде шешілетіндігі көрсетілді.

Түйін сөздер: локалды емес шекаралық есеп, Кон Лапласы, Гейзенберг группасы.

Д. Сураған

Институт математики и математического моделирования, МОН РК, Алматы, Казахстан

ОБ ОДНОЙ НЕЛОКАЛЬНОЙ КРАЕВОЙ ЗАДАЧЕ НА ГРУППЕ ГЕЙЗЕНБЕРГА

Аннотация. В статье раскрывается построение корректной краевой задачи для оператора Кон-Лапласа с нелокальным краевым условием в ограниченной гладкой области на группе Гейзенберга. А также показано, что краевая задача решается в явном виде для любой ограниченной гладкой области.

Ключевые слова: нелокальная, краевая задача, Кон Лапласа, группа Гейзенберга.

CONTENTS

<i>Saymbetov A.K., Japashov N.M., Sissenov N.K., Kuttybay N.B., Mukhametkali, B.K. Tulkibayuly Ye., Nurgaliyev M.K.</i> Development of technology and making of silicon detector structures of large size.....	15
<i>Saymbetov A.K., Japashov N.M., Sissenov N.K., Kuttybay N.B., Mukhametkali B.K., Tulkibayuly Ye., Nurgaliyev M.K.</i> Physical features of formation of silicon p-i-n detector structures.....	19
<i>Karakulova A.K., Ongeldiyeva S.M., Kuznetsova T.D.</i> Information and communication technologies at foreign language lessons as a quality factor.....	23
<i>Suragan D.</i> A nonlocal boundary value problem on the Heisenberg group.....	28
<i>Tairova G.A.</i> Linguistic peculiarities of political discourse translation.....	35
<i>Saymbetov A.K., Zhapashov N.M., Sissenov N.K., Kuttybay N.B., Mukhametkali B.K., Tulkibayuly Ye., Nurgaliyev M.K.</i> Development of high sensitive silicon strip detectors of large sizes.....	41
<i>Poleshchuk O.Kh., Dedushenko S.K., Ermakhanov M.N., Saidakhmetov P.A., Nurullaev M.A.</i> Estimations of the isomer Mössbauer shifts for tetraoxoferrates using ADF package.....	45
<i>Adizbayeva DZh., Shoybekova A.Zh.</i> Philosophical and methodological principles of pedagogy education.....	50
<i>Adizbayeva D.Zh., Shoybekova A.Zh.</i> Active personality: love, leadership and creativity as a manifestation of the phenomenon of dialog.....	55

**PUBLICATION ETHICS AND PUBLICATION MALPRACTICE
IN THE JOURNALS OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the work described has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the originality detection service Cross Check <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

www.nauka-nanrk.kz

<http://www.reports-science.kz/index.php/ru/>

Редакторы М. С. Ахметова, Д. С. Аленов, Т.А. Апендиев
Верстка на компьютере С.К. Досаевой

Подписано в печать 05.02.2016.
Формат 60х881/8. Бумага офсетная. Печать – ризограф.
3,1 п.л. Тираж 2000. Заказ 1.