

ISSN 2224-5227

2016 • 4

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
БАЯНДАМАЛАРЫ

ДОКЛАДЫ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН

REPORTS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN

ЖУРНАЛ 1944 ЖЫЛДАН ШЫҒА БАСТАҒАН
ЖУРНАЛ ИЗДАЕТСЯ С 1944 г.
PUBLISHED SINCE 1944



Бас редактор
ҚР ҰҒА академигі **М.Ж. Жұрынов**

Редакция алқасы:

хим.ғ. докторы, проф., ҚР ҰҒА академигі **Әдекенов С.М.** (бас редактордың орынбасары), эк.ғ. докторы, проф., ҚР ҰҒА академигі **Әділов Ж.М.**, мед. ғ. докторы, проф., ҚР ҰҒА академигі **Арзықұлов Ж.А.**, техн. ғ. докторы, проф., ҚР ҰҒА академигі **Бишімбаев У.К.**, а.-ш.ғ. докторы, проф., ҚР ҰҒА академигі **Есполов Т.И.**, техн. ғ. докторы, проф., ҚР ҰҒА академигі **Мұтанов Г.М.**, физ.-мат.ғ. докторы, проф., ҚР ҰҒА академигі **Өтелбаев М.О.**, пед. ғ. докторы, проф., ҚР ҰҒА академигі **Пралиев С.Ж.**, геогр.ғ. докторы, проф., ҚР ҰҒА академигі **Северский И.В.**; тарих.ғ. докторы, проф., ҚР ҰҒА академигі **Сыдықов Е.Б.**, физ.-мат.ғ. докторы, проф., ҚР ҰҒА академигі **Тәкібаев Н.Ж.**, физ.-мат.ғ. докторы, проф., ҚР ҰҒА академигі **Харин С.Н.**, тарих ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Әбүсейітова М.Х.**, экон. ғ. докторы, проф., ҰҒА корр. мүшесі **Бейсембетов И.К.**, биол. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Жамбакин К.Ж.**, тарих ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Кәрібаев Б.Б.**, мед. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Локшин В.Н.**, геол.-мин. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Өмірсеріков М.Ш.**, физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Рамазанов Т.С.**, физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Садыбеков М.А.**, хим.ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Сатаев М.И.**; ҚР ҰҒА құрметті мүшесі, а.-ш.ғ. докторы, проф. **Омбаев А.М.**

Редакция кеңесі:

Украинаның ҰҒА академигі **Гончарук В.В.** (Украина), Украинаның ҰҒА академигі **Неклюдов И.М.** (Украина), Беларусь Республикасының ҰҒА академигі **Гордиенко А.И.** (Беларусь), Молдова Республикасының ҰҒА академигі **Дука Г.** (Молдова), Тәжікстан Республикасының ҰҒА академигі **Илолов М.И.** (Тәжікстан), Қырғыз Республикасының ҰҒА академигі **Эркебаев А.Э.** (Қырғызстан), Ресей ҒА корр. мүшесі **Величкин В.И.** (Ресей Федерациясы); хим.ғ. докторы, профессор **Марек Сикорски** (Польша), тех.ғ. докторы, профессор **Потапов В.А.** (Украина), биол.ғ. докторы, профессор **Харун Парлар** (Германия), профессор **Гао Энджун** (КХР), филос. ғ. докторы, профессор **Стефано Перни** (Ұлыбритания), ғ. докторы, профессор **Богуслава Леска** (Польша), философия ғ. докторы, профессор **Полина Прокопович** (Ұлыбритания), профессор **Вуйцик Вольдемар** (Польша), профессор **Нур Изура Уздир** (Малайзия), д.х.н., профессор **Нараев В.Н.** (Ресей Федерациясы)

Главный редактор
академик НАН РК **М.Ж. Журинов**

Редакционная коллегия:

доктор хим. наук, проф., академик НАН РК **С.М. Адекенов** (заместитель главного редактора), доктор экон. наук, проф., академик НАН РК **Ж.М. Адилов**, доктор мед. наук, проф., академик НАН РК **Ж.А. Арзыкулов**, доктор техн. наук, проф., академик НАН РК **В.К. Бишимбаев**, доктор сельскохоз. наук, проф., академик НАН РК **Т.И. Есполов**, доктор техн. наук, проф., академик НАН РК **Г.М. Мутанов**, доктор физ.-мат. наук, проф., академик НАН РК **М.О. Отелбаев**, доктор пед. наук, проф., академик НАН РК **С.Ж. Пралиев**, доктор геогр. наук, проф., академик НАН РК **И.В. Северский**, доктор ист. наук, проф., академик НАН РК **Е.Б. Сыдыков**, доктор физ.-мат. наук, проф., академик НАН РК **Н.Ж. Такибаев**, доктор физ.-мат. наук, проф., академик НАН РК **С.Н. Харин**, доктор ист. наук, проф., чл.-корр. НАН РК **М.Х. Абусейтова**, доктор экон. наук, проф., чл.-корр. НАН РК **И.К. Бейсембетов**, доктор биол. наук, проф., чл.-корр. НАН РК **К.Ж. Жамбакин**, доктор ист. наук, проф., чл.-корр. НАН РК **Б.Б. Карибаев**, доктор мед. наук, проф., чл.-корр. НАН РК **В.Н. Локшин**, доктор геол.-мин. наук, проф., чл.-корр. НАН РК **М.Ш. Омирсериков**, доктор физ.-мат. наук, проф., чл.-корр. НАН РК **Т.С. Рамазанов**, доктор физ.-мат. наук, проф., чл.-корр. НАН РК **М.А. Садыбеков**, доктор хим. наук, проф., чл.-корр. НАН РК **М.И. Сатаев**; почетный член НАН РК, доктор сельскохоз. наук, проф., **А.М. Омбаев**

Редакционный совет:

академик НАН Украины **Гончарук В.В.** (Украина), академик НАН Украины **И.М. Неклюдов** (Украина), академик НАН Республики Беларусь **А.И.Гордиенко** (Беларусь), академик НАН Республики Молдова **Г. Дука** (Молдова), академик НАН Республики Таджикистан **М.И. Илолов** (Таджикистан), член-корреспондент РАН **Величкин В.И.** (Россия); академик НАН Кыргызской Республики **А.Э. Эркебаев** (Кыргызстан), д.х.н., профессор **Марек Сикорски** (Польша), д.т.н., профессор **В.А. Потапов** (Украина), д.б.н., профессор **Харун Парлар** (Германия), профессор **Гао Энджун** (КНР), доктор философии, профессор **Стефано Перни** (Великобритания), доктор наук, профессор **Богуслава Леска** (Польша), доктор философии, профессор **Полина Прокопович** (Великобритания), профессор **Вуйцик Вольдемар** (Польша), профессор **Нур Изура Уздир** (Малайзия), д.х.н., профессор **В.Н. Нараев** (Россия)

«Доклады Национальной академии наук Республики Казахстан» ISSN 2224-5227

Собственник: Республиканское общественное объединение «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №5540-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год. Тираж: 2000 экземпляров

Адрес редакции: 050010, г.Алматы, ул.Шевченко, 28, ком.218-220, тел. 272-13-19, 272-13-18

<http://nauka-nanrk.kz> reports-science.kz

Адрес типографии: ИП «Аруна», г.Алматы, ул.Муратбаева, 75

©Национальная академия наук Республики Казахстан, 2016 г.

E d i t o r i n c h i e f

M.Zh. Zhurinov, academician of NAS RK

Editorial board:

S.M. Adekenov (deputy editor in chief), Doctor of Chemistry, prof., academician of NAS RK; **Zh.M. Adilov**, Doctor of Economics, prof., academician of NAS RK; **Zh.A. Arzykulov**, Doctor of Medicine, prof., academician of NAS RK; **V.K. Bishimbayev**, Doctor of Engineering, prof., academician of NAS RK; **T.I. Yespolov**, Doctor of Agriculture, prof., academician of NAS RK; **G.M. Mutanov**, Doctor of Physics and Mathematics, prof., academician of NAS RK; **M.O. Otelbayev**, Doctor of Physics and Mathematics, prof., academician of NAS RK; **S.Zh. Praliyev**, Doctor of Education, prof., academician of NAS RK; **I.V. Seversky**, Doctor of Geography, prof., academician of NAS RK; **Ye.B. Sydykov**, Doctor of Historical Sciences, prof., academician of NAS RK; **N.Zh. Takibayev**, Doctor of Physics and Mathematics, prof., academician of NAS RK; **S.N. Kharin**, Doctor of Physics and Mathematics, prof., academician of NAS RK; **M.Kh. Abuseitova**, Doctor of Historical Sciences, prof., corr. member of NAS RK; **I.K. Beisembetov**, Doctor of Economics, prof., corr. member of NAS RK; **K.Zh. Zhambakin**, Doctor of Biological Sciences, prof., corr. member of NAS RK; **B.B. Karibayev**, Doctor of Historical Sciences, prof., corr. member of NAS RK; **V.N. Lokshin**, Doctor of Medicine, prof., corr. member of NAS RK; **M.Sh. Omirserikov**, Doctor of Geology and Mineralogy, prof., corr. member of NAS RK; **T.S. Ramazanov**, Doctor of Physics and Mathematics, prof., corr. member of NAS RK; **M.A. Sadybekov**, Doctor of Physics and Mathematics, prof., corr. member of NAS RK; **M.I. Satayev**, Doctor of Chemistry, prof., corr. member of NAS RK; **A.M. Ombayev**, Honorary Member of NAS RK, Doctor of Agriculture, prof.

Editorial staff:

V.V. Goncharuk, NAS Ukraine academician (Ukraine); **I.M. Neklyudov**, NAS Ukraine academician (Ukraine); **A.I. Gordienko**, NAS RB academician (Belarus); **G. Duca**, NAS Moldova academician (Moldova); **M.I. Iolov**, NAS Tajikistan academician (Tajikistan); **A.E. Erkebayev**, NAS Kyrgyzstan academician (Kyrgyzstan); **V.I. Velichkin**, RAS corr.member (Russia); **Marek Sikorski**, Doctor of Chemistry, prof. (Poland); **V.A. Potapov**, Doctor of Engineering, prof. (Ukraine); **Harun Parlari**, Doctor of Biological Sciences, prof. (Germany); **Gao Endzhun**, prof. (PRC); **Stefano Perni**, Doctor of Philosophy, prof. (UK); **Boguslava Leska**, dr, prof. (Poland); **Pauline Prokopovich**, Doctor of Philosophy, prof. (UK); **Wójcik Waldemar**, prof. (Poland), **Nur Izura Udzir**, prof. (Malaysia), **V.N. Narayev**, Doctor of Chemistry, prof. (Russia)

Reports of the National Academy of Sciences of the Republic of Kazakhstan.**ISSN 2224-5227**

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 5540-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 2000 copies

Editorial address: 28, Shevchenko str., of.219-220, Almaty, 050010, tel. 272-13-19, 272-13-18,

<http://nauka-nanrk.kz/> reports-science.kz

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

© National Academy of Sciences of the Republic of Kazakhstan, 2016

**REPORTS OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN**

ISSN 2224-5227

Volume 4, Number 308 (2016), 37 – 48

УДК 517.957.6

Merey Sarsengeldin, Samat Kassabek, Bizhigit Sagidolla

Department of mathematics and natural sciences, Suleyman Demirel University,
Almaty, Qaskelen, Kazakhstan

E-mail: sarsengeldin.merey@sdu.edu.kz, kassabek@gmail.com, bizhigit.sagidolla@gmail.com

**EXACT AND APPROXIMATE SOLUTIONS
OF TWO PHASE INVERSE STEFAN PROBLEM**

Abstract. The main idea is finding the exact solution of two phase inverse Stefan problem for degenerate domain with moving boundary.

Tracking answers of these questions will be organized as following. For finding analytical solution we mainly follow the method proposed by S.N. Kharin applying Faa Di Bruno's formula for Integral Error Functions. In the continuation of this section Integral Error Functions and its properties necessary for elaboration of new methods are presented.

The Stefan problem is very complicated for exact solution. Therefore we try to find an approximate solution by profile assignment of the temperature with change of differential equation by heat balance equations, which are obtained by integration with respect to x and t . For finding approximate solution we mainly follow the method proposed by S.N. Kharin applying Integral Power Balance Method.

Keywords: Integral Error Functions, Integral Power Balance Method, Two Phase Inverse Stefan Problem.

Problem formulation

In the following two sections we will deal to find the exact and approximate solutions of heat equations:

$$\frac{\partial u_1}{\partial t} = a_1^2 \frac{\partial^2 u_1}{\partial x^2}, \quad 0 < x < \alpha(t), \quad t > 0 \quad (1)$$

$$\frac{\partial u_2}{\partial t} = a_2^2 \frac{\partial^2 u_2}{\partial x^2}, \quad \alpha(t) < x < \infty, \quad t > 0 \quad (2)$$

Subjected to the following conditions:

$$u_1(0, 0) = 0 \quad (3)$$

$$u_2(x, 0) = f(x) \quad \alpha(t) < x < \infty \quad (4)$$

$$-\lambda_1 \frac{\partial u_1}{\partial x} \Big|_{x=0} = P(t), \quad t > 0 \quad (5)$$

$$u_1(\alpha(t), t) = u_2(\alpha(t), t) = u_m \quad (6)$$

The Stefan's condition:

$$-\lambda_1 \frac{\partial u_1}{\partial x} \Big|_{x=\alpha(t)} = -\lambda_2 \frac{\partial u_2}{\partial x} \Big|_{x=\alpha(t)} + L\gamma \frac{d(\alpha(t))}{dt} \quad (7)$$

$$u_2 \Big|_{x=\infty} = 0 \quad (8)$$

It is necessary to find temperature distribution $u_1(x, t)$ and $u_2(x, t)$ also it is required to reconstruct the boundary function $P(t)$ if the free boundary $\alpha(t)$ is given. Such problem is called two phase inverse Stefan Problem. The heat spread in the solid is negligible because of the physical properties of contact material. This condition is valid for refractory metals like wolfram. After getting the solutions we try to discuss given both solutions.

I. EXACT SOLUTION OF TWO PHASE INVERSE STEFAN PROBLEM

1.1. Introduction

The first analytical solution of two phase inverse Stefan problem, which describes the dynamics of soil freezing has been published by Lamé and Clayperon.

Despite the quite extensive list of problems in literature which lead to the necessity to solve Stefan type problems see: e.g., [1–9] and a long bibliography [9] on methods for solving these problems lead to additional difficulties which occur due to the degeneracy of domains. In some specific cases particularly for free moving boundaries it is possible to construct Heat potentials and a problem can be reduced to the system of integral equations [2], [3], however in the case of degeneracy, singularity in integral equations occur, and method of successive approximations is inapplicable in general. Moreover, the use of numerical methods is problematic when the number of parameters is great. Therefore, development of new analytical methods is very important especially for various applications because it enables one to analyze an interrelationship of different input parameters and their influence on the dynamics of investigating phenomena.

As for applications: a wide range of electric contact phenomena, in particular, the phenomena occurring at the interaction of electrical arc with electrode can be described in dynamic use of the presented method see e.g., [9] for very short arc duration (nanosecond diapason), when experimental investigation is very difficult. In this study we will try to find solution of two Phase Inverse Stefan problem for degenerate domain with moving boundary.

1.2. Problem Solution

Let us consider this problem at the suggestion that $\alpha(t) = \sum_{n=1}^{\infty} \alpha_n t^{n/2}$ is given.

The unknown functions are $u_1(x, t), u_2(x, t), P(t)$ we represent in the form of Heat polynomials and Integral Error Functions:

$$u_1(x, t) = \sum_{n=0}^{\infty} \left[(2a_1 \sqrt{t})^{2n} A_{2n} i^{2n} \operatorname{erfc} \left(\frac{x}{2a_1 \sqrt{t}} \right) + (2a_1 \sqrt{t})^{2n+1} A_{2n+1} i^{2n+1} \operatorname{erfc} \left(-\frac{x}{2a_1 \sqrt{t}} \right) \right] \quad (9)$$

$$u_2(x, t) = \sum_{n=0}^{\infty} \left[(2a_2 \sqrt{t})^{2n} B_{2n} i^{2n} \operatorname{erfc} \left(\frac{x}{2a_2 \sqrt{t}} \right) + (2a_2 \sqrt{t})^{2n+1} B_{2n+1} i^{2n+1} \operatorname{erfc} \left(-\frac{x}{2a_2 \sqrt{t}} \right) \right] \quad (10)$$

$$P(t) = \sum_{n=0}^{\infty} P_n t^{\frac{n}{2}} \quad (11)$$

Where the coefficients $A_{2k}, A_{2k+1}, B_{2k}, B_{2k+1}, P(t)$ should be found satisfying the boundary conditions.

Let us satisfy first the initial condition (4). It is clear that:

$$u_2(x, 0) = \lim_{t \rightarrow 0} u_2(x, t) = \sum_{n=0}^{\infty} \left[\lim_{t \rightarrow 0} \left(2a_2 \sqrt{t} \right)^{2n} B_{2n} i^{2n} \operatorname{erfc} \left(\frac{x}{2a_2 \sqrt{t}} \right) + \lim_{t \rightarrow 0} \left(2a_2 \sqrt{t} \right)^{2n+1} B_{2n+1} i^{2n+1} \operatorname{erfc} \left(-\frac{x}{2a_2 \sqrt{t}} \right) \right] \tag{12}$$

by properties of Integral Error Functions:

$$\lim_{x \rightarrow \infty} \frac{i^n \operatorname{erfc}(x)}{x^n} = 0$$

by using L'Hospital Rule to find limit:

$$\lim_{x \rightarrow \infty} \frac{i^n \operatorname{erfc}(-x)}{x^n} = \lim_{x \rightarrow \infty} \frac{i^{n-1} \operatorname{erfc}(-x)}{nx^{n-1}} = \lim_{x \rightarrow \infty} \frac{i^{n-2} \operatorname{erfc}(-x)}{n(n-1)x^{n-2}} = \dots = \lim_{x \rightarrow \infty} \frac{\operatorname{erfc}(-x)}{n!} = \frac{2}{n!}$$

So therefore,

$$u_2(x, 0) = \sum_{n=0}^{\infty} B_{2n+1} \frac{2}{(2n+1)!} x^{2n+1} = f(x) \tag{13}$$

If we expand $f(x)$ into Taylor series:

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(2n+1)}(0)}{(2n+1)!} x^{2n+1} \tag{14}$$

$$\sum_{n=0}^{\infty} B_{2n+1} \frac{2}{(2n+1)!} x^{2n+1} = \sum_{n=0}^{\infty} \frac{f^{(2n+1)}(0)}{(2n+1)!} x^{2n+1} \quad B_{2n+1} = \frac{1}{2} f^{(2n+1)}(0), \quad (n = 0, 1, 2, \dots) \tag{15}$$

From the condition (6) for $u_1(\alpha(t), t) = u_m$:

$$u_2(x, t) = \sum_{n=0}^{\infty} \left[\left(2a_2 \sqrt{t} \right)^{2n} B_{2n} i^{2n} \operatorname{erfc} \left(\frac{x}{2a_2 \sqrt{t}} \right) + \left(2a_2 \sqrt{t} \right)^{2n+1} B_{2n+1} i^{2n+1} \operatorname{erfc} \left(-\frac{x}{2a_2 \sqrt{t}} \right) \right] \Big|_{x=\alpha(t)} = u_m$$

The main problem is to find the remaining unknown coefficients we have, also if $\sqrt{t} = \tau$, then our expression will get:

$$\begin{aligned} \frac{\partial^{2k}}{\partial \tau^{2k}} \left(\sum_{n=0}^{\infty} \left[\left(2a_2 \tau \right)^{2n} B_{2n} i^{2n} \operatorname{erfc} \left(\frac{\alpha(\tau^2)}{2a_2 \tau} \right) + \left(2a_2 \tau \right)^{2n+1} B_{2n+1} i^{2n+1} \operatorname{erfc} \left(-\frac{\alpha(\tau^2)}{2a_2 \tau} \right) \right] \right) &= \frac{\partial^{2k}}{\partial \tau^{2k}} (u_m) = 0 \\ \sum_{n=0}^{\infty} \sum_{l=0}^{2k} \left(2a_2 \right)^{2n} B_{2n} \binom{2k}{l} \frac{(2n)!}{(2n-2k+l)!} \cdot \tau^{2n-2k+l} \cdot \frac{\partial^{2k}}{\partial \tau^{2k}} \left(i^{2n} \operatorname{erfc} \left(\frac{\alpha(\tau^2)}{2a_2 \tau} \right) \right) &+ \\ + \left(2a_2 \right)^{2n+1} B_{2n+1} \binom{2k}{l} \frac{(2n)!}{(2n+1-2k+l)!} \cdot \tau^{2n+1-2k+l} \cdot \frac{\partial^{2k}}{\partial \tau^{2k}} \left(i^{2n+1} \operatorname{erfc} \left(-\frac{\alpha(\tau^2)}{2a_2 \tau} \right) \right) &= 0 \end{aligned}$$

After taking both parts 2k-derivatives we can see following expression:

$$\sum_{n=0}^{\infty} \left(2a_2 \right)^{2n} B_{2n} \frac{(2k)!}{(2k-2n)!} \frac{\partial^{2k-2n}}{\partial \tau^{2k-2n}} \left(i^{2n} \operatorname{erfc} \left(\frac{\alpha(\tau^2)}{2a_2 \tau} \right) \right) \Big|_{\tau=0} +$$

$$+(2a_2)^{2n+1} B_{2n+1} \frac{(2k)!}{(2k-2n-1)!} \frac{\partial^{2k-2n-1}}{\partial \tau^{2k-2n-1}} \left(i^{2n+1} \operatorname{erfc} \left(-\frac{\alpha(\tau^2)}{2a_2\tau} \right) \right) \Big|_{\tau=0} = 0$$

For this purpose we use the Faa Di Bruno formula (Arbogast formula) for a derivative of a composite function:

$$\begin{aligned} \frac{\partial^{2k-2n}}{\partial \tau^{2k-2n}} \left(i^{2n} \operatorname{erfc} \left(\frac{\alpha(\tau^2)}{2a_2\tau} \right) \right) &= \sum_{j=0}^{2k-2n} i^{2n} \operatorname{erfc}^{(j)}(\delta) \beta_{2k-2n,j}(\delta', \delta'', \dots, \delta^{2k-2n-j}) \\ i^{2n} \operatorname{erfc}^{(j)}(\delta) &= \frac{d^j}{d\tau^j} \left(i^{2n} \operatorname{erfc}(\delta) \right) \Big|_{\tau=0} = (-1)^{j-2n-1} \frac{2}{\sqrt{\pi}} H_{j-2n-1}(\delta) \exp(-\delta^2) \\ \frac{\partial^{2k-2n-1}}{\partial \tau^{2k-2n-1}} \left(i^{2n+1} \operatorname{erfc} \left(\frac{\alpha(\tau^2)}{2a_2\tau} \right) \right) &= \sum_{j=0}^{2k-2n-1} i^{2n+1} \operatorname{erfc}^{(j)}(\delta) \beta_{2k-2n-1,j}(\delta', \delta'', \dots, \delta^{2k-2n-1-j}) \\ i^{2n+1} \operatorname{erfc}^{(j)}(\delta) &= \frac{d^j}{d\tau^j} \left(i^{2n+1} \operatorname{erfc}(\delta) \right) \Big|_{\tau=0} = (-1)^{j-2n-2} \frac{2}{\sqrt{\pi}} H_{j-2n-2}(\delta) \exp(-\delta^2) \end{aligned}$$

Let us denote our boundaries in the following form to get rid of the singularity:

$$\begin{aligned} \gamma(t) &= \alpha(t) - \alpha_0 = \gamma_1 t^{1/2} + \gamma_2 t + \dots = \sum_{n=0}^{\infty} \gamma_{n+1} t^{(n+1)/2} \\ \alpha(\tau^2) &= \gamma_1 \tau + \gamma_2 \tau^2 + \dots + \gamma_{n+1} \tau^{n+1} = \sum_{n=0}^{\infty} \gamma_{n+1} \tau^{n+1} \end{aligned}$$

Then we can easily find our boundaries:

$$\begin{aligned} \delta \Big|_{\tau=0} &= \frac{\sum_{n=0}^{\infty} \gamma_{n+1} \tau^{n+1}}{2a_2\tau} \Big|_{\tau=0} = \frac{\gamma_1}{2a_2} \\ \delta \Big|_{\tau=0} &= -\frac{\sum_{n=0}^{\infty} \gamma_{n+1} \tau^{n+1}}{2a_2\tau} \Big|_{\tau=0} = -\frac{\gamma_1}{2a_2} \end{aligned}$$

Finally, we get the following sum:

$$\begin{aligned} \sum_{n=0}^{\infty} B_{2n} \frac{(2a_2)^{2n} (2k)!}{(2k-2n)!} \sum_{j=0}^{2k-2n} (-1)^{j-2n-1} \frac{2}{\sqrt{\pi}} H_{j-2n-1} \left(\frac{\gamma_1}{2a_2} \right) \exp \left(-\frac{\gamma_1^2}{4a_2^2} \right) \beta_{2k-2n,j}(\delta', \delta'', \dots, \delta^{2k-2n-j}) + \\ + \sum_{n=0}^{\infty} B_{2n+1} \frac{(2a_2)^{2n+1} (2k)!}{(2k-2n-1)!} \sum_{j=0}^{2k-2n-1} (-1)^{j-2n} \frac{2}{\sqrt{\pi}} H_{j-2n-2} \left(-\frac{\gamma_1}{2a_2} \right) \exp \left(-\frac{\gamma_1^2}{4a_2^2} \right) \beta_{2k-2n-1,j}(\delta', \delta'', \dots, \delta^{2k-2n-1-j}) = 0 \end{aligned}$$

Here let us denote the known coefficients in the following form:

$$K_{n,1} = \sum_{n=0}^{\infty} \frac{(2a_2)^{2n} (2k)!^{2k-2n}}{(2k-2n)!} \sum_{j=0}^{2k-2n} (-1)^{j-2n-1} \frac{2}{\sqrt{\pi}} H_{j-2n-1} \left(\frac{\gamma_1}{2a_2} \right) \exp \left(-\frac{\gamma_1^2}{4a_2^2} \right) \beta_{2k-2n,j} (\delta', \delta'', \dots, \delta^{2k-2n-j})$$

$$K_{n,2} = \sum_{n=0}^{\infty} \frac{(2a_2)^{2n+1} (2k)!^{2k-2n-1}}{(2k-2n-1)!} \sum_{j=0}^{2k-2n-1} (-1)^{j-2n} \frac{2}{\sqrt{\pi}} H_{j-2n-2} \left(-\frac{\gamma_1}{2a_2} \right) \exp \left(-\frac{\gamma_1^2}{4a_2^2} \right) \beta_{2k-2n-1,j} (\delta', \delta'', \dots, \delta^{2k-2n-1-j})$$

After we can easily find our unknown coefficient:

$$B_{2n} \cdot K_{n,1} = B_{2n+1} \cdot K_{n,2}$$

$$B_{2n} = \frac{B_{2n+1} \cdot K_{n,2}}{K_{n,1}} = \frac{f^{(2n+1)}(0) \cdot K_{n,2}}{2K_{n,1}} \tag{16}$$

From the condition (6) for $u_1(x, t)$ will be the same situation for $u_1(\alpha(t), t) = u_m$:

$$A_{2n} = \frac{A_{2n+1} \cdot K_{n,2}}{K_{n,1}} \tag{17}$$

By using the condition (7):

$$dt = 2\tau d\tau$$

$$\left(\frac{L\gamma}{2\tau} \frac{d\alpha(\tau)}{d\tau} \right) \Big|_{\tau=0}^{(k)} = \left(\frac{L\gamma}{2} \sum_{n=0}^{\infty} \frac{\alpha^{(n)}(0)}{(2n)!} \tau^{2n-1} \right) \Big|_{\tau=0}^{(k)}$$

It is easy to see that $k = 2n - 1$, therefore:

$$\frac{L\gamma}{2} \sum_{n=0}^{\infty} \frac{\alpha^{(n)}(0)}{(2n)!} (\tau^{2n-1}) \Big|_{\tau=0}^{(k)} = \frac{L\gamma}{2} \sum_{n=0}^{\infty} \frac{\alpha^{(n)}(0)}{2n}$$

If we will take $2n - 1$ derivatives in both parts by using Leibniz and Faa di Bruno formulas:

$$\left(-\lambda_1 \frac{\partial u_1}{\partial x} \Big|_{x=\alpha(\tau^2)} \right) \Big|_{\tau=0}^{(2n-1)} = \left(-\lambda_2 \frac{\partial u_2}{\partial x} \Big|_{x=\alpha(\tau^2)} \right) \Big|_{\tau=0}^{(2n-1)} + \frac{L\gamma}{4} \sum_{n=0}^{\infty} \frac{\alpha^{(n)}(0)}{n}$$

$$\left[\lambda_1 \sum_{n=0}^{\infty} (2a_1)^{2n-1} A_{2n} (2n-1)! \left(i^{2n-1} \operatorname{erfc} \left(\frac{\alpha(\tau^2)}{2a_1\tau} \right) \right) \Big|_{\tau=0} - \right.$$

$$\left. -\lambda_1 \sum_{n=0}^{\infty} (2a_1)^{2n} A_{2n+1} \cdot \tau \cdot (2n-1)! \left(i^{2n} \operatorname{erfc} \left(-\frac{\alpha(\tau^2)}{2a_1\tau} \right) \right) \Big|_{\tau=0} \right] =$$

$$\lambda_1 \sum_{n=0}^{\infty} (2a_1)^{2n-1} A_{2n} (2n-1)! \left(i^{2n-1} \operatorname{erfc} \left(\frac{\gamma_1 + \gamma_2\tau + \dots}{2a_1} \right) \right) \Big|_{\tau=0} = A_{2n} \sum_{n=0}^{\infty} \lambda_1 (2a_1)^{2n-1} (2n-1)! \left(i^{2n-1} \operatorname{erfc} \left(\frac{\gamma_1}{2a_1} \right) \right)$$

$$\sum_{n=0}^{\infty} A_{2n} C_{n,1} = \sum_{n=0}^{\infty} B_{2n} C_{n,2} + \frac{L\gamma}{4} \sum_{n=0}^{\infty} \frac{\alpha^{(n)}(0)}{n} \tag{18}$$

where

$$C_{n,1} = \lambda_1 (2a_1)^{2n-1} (2n-1)! i^{2n-1} \operatorname{erfc} \left(\frac{\gamma_1}{2a_1} \right)$$

$$C_{n,2} = \lambda_2 (2a_2)^{2n-1} (2n-1)! i^{2n-1} \operatorname{erfc} \left(\frac{\gamma_1}{2a_2} \right)$$

From (17) and (18) we can easily find two unknowns in the following form:

$$A_{2n} = \frac{4n \cdot C_{n,2} \cdot f^{(2n+1)}(0) \cdot K_{n,2} + L\gamma \cdot \alpha^{(n)}(0) K_{n,1}}{8n \cdot K_{n,1} \cdot C_{n,1}} \quad (19)$$

$$A_{2n+1} = \frac{4n \cdot C_{n,2} \cdot f^{(2n+1)}(0) \cdot K_{n,2} + L\gamma \cdot \alpha^{(n)}(0) K_{n,1}}{8n \cdot K_{n,2} \cdot C_{n,1}} \quad (20)$$

By using the last condition (5) we get:

$$\lambda_1 \sum_{n=0}^{\infty} (2a_1)^{2n-1} \left[A_{2n}(\tau)^{2n-1} i^{2n-1} \operatorname{erfc}(0) - A_{2n+1}(\tau)^{2n} i^{2n} \operatorname{erfc}(0) \right] = \sum_{n=0}^{\infty} P_n \tau^n \quad (21)$$

The main problem is to find the remaining unknown coefficient we have:

$$\begin{aligned} \frac{\partial^k}{\partial \tau^k} \left(\lambda_1 \sum_{n=0}^{\infty} \left[(2a_1 \tau)^{2n-1} A_{2n} i^{2n-1} \operatorname{erfc} \left(\frac{x}{2a_1 \tau} \right) + (2a_1 \tau)^{2n} A_{2n+1} i^{2n} \operatorname{erfc} \left(-\frac{x}{2a_1 \tau} \right) \right] \right) &= \frac{\partial^k}{\partial \tau^k} \left(\sum_{n=0}^{\infty} P_n \tau^n \right) \\ \lambda_1 \sum_{n=0}^{\infty} \sum_{l=0}^k \left[(2a_1)^{2n-1} A_{2n} \binom{k}{l} \frac{(2n-1)!}{(2n-1-k+l)!} \cdot \tau^{2n-1-k+l} \cdot \frac{\partial^k}{\partial \tau^k} \left(i^{2n-1} \operatorname{erfc} \left(\frac{x}{2a_1 \tau} \right) \right) - \right. \\ \left. - (2a_1)^{2n} A_{2n+1} \binom{k}{l} \frac{(2n)!}{(2n-k+l)!} \cdot \tau^{2n-k+l} \cdot \frac{\partial^k}{\partial \tau^k} \left(i^{2n} \operatorname{erfc} \left(-\frac{x}{2a_1 \tau} \right) \right) \right] &= \frac{\partial^k}{\partial \tau^k} \left(\sum_{n=0}^{\infty} P_n \tau^n \right) \end{aligned}$$

After taking both parts k-derivatives we can see following expression:

$$\begin{aligned} \lambda_1 \left[\sum_{n=0}^{\infty} (2a_1)^{2n-1} A_{2n} \frac{(k)!}{(k-2n+1)!} \frac{\partial^{k-2n+1}}{\partial \tau^{k-2n+1}} \left(i^{2n-1} \operatorname{erfc} \left(\frac{x}{2a_1 \tau} \right) \right) \right]_{\tau=0} - \\ - (2a_1)^{2n} A_{2n+1} \frac{(k)!}{(k-2n)!} \frac{\partial^{k-2n}}{\partial \tau^{k-2n}} \left(i^{2n} \operatorname{erfc} \left(-\frac{x}{2a_2 \tau} \right) \right) \Big|_{\tau=0} = \left(\sum_{n=0}^{\infty} P_n \frac{k!}{(k-n)!} \right) \end{aligned}$$

For this purpose we use the Faa Di Bruno formula (Arbogast formula) for a derivative of a composite function:

$$\begin{aligned} \frac{\partial^{k-2n+1}}{\partial \tau^{k-2n+1}} \left(i^{2n-1} \operatorname{erfc} \left(\frac{x}{2a_2 \tau} \right) \right) &= \sum_{j=0}^{k-2n+1} i^{2n-1} \operatorname{erfc}^{(j)}(\delta) \beta_{k-2n+1,j}(\delta', \delta'', \dots, \delta^{k-2n+1-j}) \\ i^{2n-1} \operatorname{erfc}^{(j)}(\delta) &= \frac{d^j}{d\tau^j} \left(i^{2n-1} \operatorname{erfc}(\delta) \right) \Big|_{\tau=0} = (-1)^{j-2n} \frac{2}{\sqrt{\pi}} H_{j-2n}(\delta) \exp(-\delta^2) \end{aligned}$$

$$\delta \Big|_{x=0} = \frac{x}{2a_2\tau} \Big|_{x=0} = 0 \quad \exp(0) = 1$$

$$\frac{\partial^{k-2n}}{\partial \tau^{k-2n}} \left(i^{2n} \operatorname{erfc} \left(-\frac{x}{2a_2\tau} \right) \right) = \sum_{j=0}^{k-2n} i^{2n} \operatorname{erfc}^{(j)}(\delta) \beta_{k-2n,j}(\delta', \delta'', \dots, \delta^{k-2n-j})$$

$$i^{2n} \operatorname{erfc}^{(j)}(\delta) = \frac{d^j}{d\tau^j} \left(i^{2n} \operatorname{erfc}(\delta) \right) \Big|_{\tau=0} = (-1)^{j-2n-1} \frac{2}{\sqrt{\pi}} H_{j-2n-1}(\delta) \exp(-\delta^2)$$

Finally, we get following sum:

$$\lambda_1 \left[\sum_{n=0}^{\infty} A_{2n} \frac{(2a_1)^{2n-1} (k)!}{(k-2n+1)!} \sum_{j=0}^{k-2n+1} (-1)^{j-2n} \frac{2}{\sqrt{\pi}} H_{j-2n}(0) \beta_{k-2n+1,j}(\delta', \delta'', \dots, \delta^{k-2n+1-j}) - \right.$$

$$\left. - \sum_{n=0}^{\infty} A_{2n+1} \frac{(2a_1)^{2n} (k)!}{(k-2n)!} \sum_{j=0}^{k-2n} (-1)^{j-2n-1} \frac{2}{\sqrt{\pi}} H_{j-2n-1}(0) \beta_{k-2n,j}(\delta', \delta'', \dots, \delta^{k-2n-j}) \right] = \sum_{n=0}^{\infty} P_n \frac{k!}{(k-n)!}$$

After we can easily find our unknown coefficient:

$$A_{2n} \cdot K_{n,3} - A_{2n+1} \cdot K_{n,4} = P_n \tag{22}$$

1.3. Conclusion

Main result namely coefficients of function A_{2k} , A_{2k+1} (9), B_{2k} , B_{2k+1} (10) and the function $P(t)$ are obtained from (19), (20), (16), (15) and (21-22) respectively. It worth nothing that deviation of solution can provide maximum principle. Moreover if convergence of (9) and (10) proved, wide class of heat equations that describe diverse phenomena can be solved both analytically and numerically.

II. APPROXIMATE SOLUTION OF TWO PHASE INVERSE STEFAN PROBLEM

2.1. Introduction

Exact solutions of heat conduction problems are rather cumbersome and time-consuming. In addition, they are practically absent in the problems of the radial heat flux in spherical coordinates with the change in the aggregate state [11], [12]. Therefore, to solve practical problems are commonly used charts, obtained by numerical or approximate methods [13]. One of the approximate analytical methods is an integral method of heat balance, which primarily attracted to its physical clarity, simplicity and precision of the results is high enough that demonstrates T. Goodman [14] on numerous examples. The main difficulty to be faced when using the integral method of heat balance is the setting right temperature profile, which according to T. Goodman, significantly affects the accuracy of the results.

There are several approaches in selecting the temperature profiles. In [15] A.I. Veinik offers to use temperature profiles in the form of ordinary polynomials for problems of any geometry, which should simplify the solution of the problem.

In this study we will try to find approximate solution of two Phase Inverse Stefan problem for degenerate domain with moving boundary.

2.2. Problem Solution

Let us consider this problem at the suggestion that $\alpha(t) = \sum_{n=1}^{\infty} \alpha_n t^{n/2}$ is given.

To find the unknown $u_2(x, t)$, we identify the temperature profile corresponding following conditions:

$$u_2(\alpha(t), t) = u_m \quad (23)$$

$$u_2|_{\beta(t)} = 0 \quad (24)$$

$$\frac{\partial u_2}{\partial x} \Big|_{x=\beta(t)} = 0 \quad (25)$$

$$u_2|_{x=\infty} = 0 \quad (26)$$

Therefore the temperature profile will be:

$$u_2(x, t) = \begin{cases} u_m \left[\frac{x - \beta(t)}{\alpha(t) - \beta(t)} \right]^2, & \alpha(t) < x < \beta(t) \\ 0, & x > \beta(t) \end{cases} \quad (27)$$

Here we need to find the $\beta(t)$, where $\beta(0) = 0$ by using the equation (2) to use the integral of power balance:

$$\int_{\alpha(t)}^{\beta(t)} \frac{\partial u_2}{\partial t} dx = a_2^2 \frac{\partial u_2}{\partial x} \Big|_{\alpha(t)}^{\beta(t)} \quad (28)$$

$$\int_{\alpha(t)}^{\beta(t)} \frac{\partial u_2}{\partial t} dx = a_2^2 \left[\frac{\partial u_2}{\partial x} \Big|_{x=\beta(t)} - \frac{\partial u_2}{\partial x} \Big|_{x=\alpha(t)} \right]$$

$$\int_{\alpha(t)}^{\beta(t)} \frac{\partial u_2}{\partial t} dx = a_2^2 \left[0 - 2u_m \left[\frac{x - \beta(t)}{\alpha(t) - \beta(t)} \right] \left(\frac{1}{\alpha(t) - \beta(t)} \right) \right]_{x=\alpha(t)}$$

$$\int_{\alpha(t)}^{\beta(t)} \frac{\partial u_2}{\partial t} dx = -a_2^2 \frac{2u_m}{\alpha(t) - \beta(t)}$$

For the right side we use the Leibniz integral rule, also differentiation under the integral sign:

$$\frac{d}{dx} \left(\int_{\alpha(x)}^{b(x)} f(x, t) dt \right) = f(x, b(x)) \cdot b'(x) - f(x, \alpha(x)) \cdot \alpha'(x) + \int_{\alpha(x)}^{b(x)} \frac{\partial}{\partial x} f(x, t) dt$$

Therefore,

$$\int_{\alpha(t)}^{\beta(t)} \frac{\partial}{\partial t} f(x, t) dx = \frac{d}{dt} \left(\int_{\alpha(x)}^{\beta(x)} f(x, t) dx \right) - f(\beta(t), t) \cdot \beta'(t) + f(\alpha(t), t) \cdot \alpha'(t) \quad (29)$$

By using Leibniz’s method we get:


$$\left[\frac{d}{dt} \left(\int_{\alpha(t)}^{\beta(t)} u_2 dx \right) - u_2 |_{x=\beta(t)} \frac{d\beta(t)}{dt} + u_2 |_{x=\alpha(t)} \frac{d\alpha(t)}{dt} \right] = -a_2^2 \frac{2u_m}{\alpha(t) - \beta(t)}$$

The solution of given differential equation has been shown in the paper [24].

$$\frac{d\beta(t)}{dt} (\beta(t) - \alpha(t)) + 2 \frac{d\alpha(t)}{dt} (\beta(t) - \alpha(t)) - 6a_2^2 = 0 \tag{30}$$


To solve the equation (30) we have used the mathematical program Mathcad 13. To find the solution we considered three particular cases and got following graphs:

$$\frac{d}{dt} y(t)(y(t) - \alpha(t)) + 2 \frac{d\alpha(t)}{dt} (y(t) - \alpha(t)) - 6a_2^2 = 0 \tag{31}$$




Equation (31) have been plotted in the following way for the first particular case when $t = 100$:

Figure 1 - $\beta(t)$ at $t = 100$



Equation (31) have been plotted in the following way for the second particular case when $t = 300$:

Figure 2 - $\beta(t)$ at $t = 300$



Equation (31) have been plotted in the following way for the third particular case when $t = 500$:

Figure 3 - $\beta(t)$ at $t = 500$

This given plots approximately show the value of $\beta(t)$. Therefore, we can consider that we know the value of $\beta(t)$ and consider other cases.

For $u_1(x, t)$ we have the temperature profile $A(t)x^2 + B(t)x + C(t)$ as it is linear. We will use the conditions (5),(6) for $u_1(x, t)$, (7) and (1):

$$u_1(x, t) = A(t)x^2 + B(t)x + C(t), \quad 0 < x < \alpha(t)$$

By using the condition (5) we get:

$$\left. \frac{\partial u_1}{\partial x} \right|_{x=0} = B(t) = -\frac{P(t)}{\lambda_1} \quad (32)$$

By using the condition (6) for $u_1(x, t)$ we have:

$$u_1(\alpha(t), t) = A(t)\alpha^2(t) + B(t)\alpha(t) + C(t) = u_m \quad (33)$$

By using the Stefan condition (7) we get:

$$-\lambda_1(2A(t)\alpha(t) + B(t)) = -\lambda_2 \left(\frac{2u_m}{\alpha(t) - \beta(t)} \right) + L\gamma \frac{d\alpha(t)}{dt}$$

If we equalize the right side of the equation above to $Q(t)$, we will have following expression:

$$2A(t)\alpha(t) + B(t) = -\frac{Q(t)}{\lambda_1} \quad (34)$$

Let us use the last equation (1) and get an integral of power balance:

$$\int_0^{\alpha(t)} \frac{\partial^2 u_1}{\partial x^2} dx = \left. \frac{\partial u_1}{\partial x} \right|_0^{\alpha(t)}$$

$$a_1^2 \left[\frac{\partial u_1}{\partial x} \Big|_{x=\alpha(t)} - \frac{\partial u_1}{\partial x} \Big|_{x=0} \right] = \int_0^{\alpha(t)} \frac{\partial u_1}{\partial t} dx$$

$$a_1^2 [2A(t) \cdot \alpha(t) + B(t) - B(t)] = a_1^2 [2A(t) \cdot \alpha(t)] = \int_0^{\alpha(t)} \frac{\partial u_1}{\partial t} dx$$

The solutions has been shown in the paper [10].

$$A'(t) [2\alpha^2(t) + 4\alpha'(t) \cdot \alpha(t)] + A(t) [4\alpha'(t) + 4(\alpha''(t) \cdot \alpha(t) + (\alpha'(t))^2) - 12a_1^2] +$$

$$+ \left[\frac{d}{dt} \left[\frac{3\alpha'(t)Q(t)}{\lambda_1} + \frac{3\alpha(t)Q'(t)}{\lambda_1} \right] - u_m \frac{\alpha'(t)}{\alpha(t)} \right] = 0 \quad (35)$$

If we denote the functions that is multiplied to $A'(t)$, $A(t)$ and some functions by $Z_1(t)$, $Z_2(t)$ and $Z_3(t)$ respectively, then we get:

$$A'(t) \cdot Z_1(t) + A(t) \cdot Z_2(t) + Z_3(t) = 0$$

We can easily rewrite our differential equation in the following form:

$$y'(x) + y(x) \cdot S(x) + T(x) = 0$$

By solving our differential equation by Integrating Factor method that has been shown in the paper [10], we can show the solution of givendifferential equation:

$$A(t) = -e^{-\int S(t)dt} \cdot \left(\int T(t) \cdot e^{\int S(t)dt} dt \right) + c \cdot e^{-\int S(t)dt} \quad (36)$$

where $S(t) = Z_2(t) / Z_1(t)$ and $T(t) = Z_3(t) / Z_1(t)$. Let us denote the right side of above expression as $K(t)$, therefore we can show the temperature profile for $u_1(x, t)$ and reconstruct the boundary function $P(t)$:

$$u_1(x, t) = K(t)x^2 - \left[\frac{Q(t)}{\lambda_1} + 2K(t)\alpha(t) \right] x + \left[u_m + K(t)\alpha^2(t) + \frac{Q(t)\alpha(t)}{\lambda_1} \right] \quad (37)$$

$$P(t) = Q(t) + \frac{2K(t)\alpha(t)}{\lambda_1} \quad (38)$$

2.4. Conclusion

In this chapter for solving problems of heat conduction in spherical geometry by integral method of power balance for the first time evaluated the effectiveness of the temperature type profiles: $T(r, t) = \text{polynomial}$ and $T(r, t) = \text{polynomial} / r$ with polynomials of degree n. The results show that in dealing with problems of thermal conductivity for different geometry approximate integral method of power balance is not always possible to achieve sufficient accuracy when using temperature profiles of the same type.

REFERENCES

- [1] V. Alexiades and A. D. Solomon, *Mathematical Modeling of Melting and Freezing Processes*, Taylor and Francis, Washington, DC, **1993**.
 [2] M. Grassi, G. Grassi, *Mathematical modelling and controlled drug delivery: matrix systems*, *Current Drug Delivery 2* (2005) 97-116.

- [3] Kharin S.N., Sarsengeldin M. Influence of Contact Materials on Phenomena in a Short Electrical Arc. TransTechPublications. Switzerland.
- [4] D.A. Tarzia, *A bibliography on moving-free boundary problems for the heat-diffusion equation. The Stefan and related problems*, 2000.
- [5] A.N. Tikhonov, A.A. Samarski, Equations of Mathematical Physics.
- [6] A. Friedman, *Free boundary problems for parabolic equations I. Melting of solids*, J. Math. Mech., 8 (1959) , pp. 499–517.
- [7] L. I. Rubinstein, *The Stefan Problem*, Transl. Math. Monogr. 1971.
- [8] Sarsengeldin M.M. “Solution Of Heat Equation With Moving Boundaries Obtained By Hartree Functions”, International scientific conference.
- [9] S.N. Kharin, M.M. Sarsengeldin, A. Temirkul, “Analytical Solution Of The Third Boundary-Value Problem For The Heat Equation By IEF Method”.
- [10] M.M. Sarsengeldin, S. Kassabek, B.M. Sagidolla, “Approximate Solution of Two phase Inverse Stefan problem” (accepted), 2016.
- [11] Carslow G., Eger D. Thermal conductivity of solids, 1964. – 488 p.
- [12] Lykov A.V. Теория теплопроводности, 1967. – 600 p.
- [13] Pehovich A.I., Zhidkih Расчеты теплового режима твердых тел, 1976.
- [14] Goodman T. Application of integral method in nonlinear problems of non-stationary heat exchange, 1967. p. 47–96.
- [15] Veinik A.I. Приближенный расчет процессов теплопроводности, 1959.

М.М. Сарсенгельдин, С. Қасабек, Б.М. Сағидолла

Сүлеймен Демирел атындағы университет, Қаскелең, Қазақстан

НАҚТЫ ЖӘНЕ ЖУЫҚТАЛҒАН ЕКІ ФАЗАЛЫ КЕРІ СТЕФАН МӘСЕЛЕЛЕРІНІҢ ШЕШІМІ

Аннотация. Берілген жұмыстың негізгі идеясы, қозғалатын шекаралары бар нұксанды өңір үшін екі фазалық кері Стефан мәселенің нақты шешімдерін табу болып табылады.

Төменде осы мәселелерді жауап қадағалау ұйымдастырылады. Аналитикалық шешім табу үшін біз негізінен С.Н. Харин жұмыстарында ұсынған әдістері Фаа Ди Бруно формуласын интегралды қателіктер функцияларына қолданамыз. Интегралды қателіктер функциясы және оның жаңа әдістерінің қасиеттері де анық көрінген.

Стефан мәселесі нақты шешім үшін күрделі болып табылады. Сондықтан, біз шамамен шешімін табу үшін, X және t арқылы интегралдап алынған дифференциалдық теңдеулерге температуралық профильдер тағайындаймыз. Бұл шамамен шешімін табу үшін С.Н. Харинның ұсынған әдістерін қолдана отырып, интегралды жылулық тепе-теңдік әдісін қолдандық.

Тірек сөздер: Интегралды функциялар қателіктер әдісі, Интегралды жылулық тепе-теңдік әдісі, Екі фазалы кері Стефан мәселесі.

М.М. Сарсенгельдин, С. Қасабек, Б.М. Сағидолла

Университет им. Сулеймана Демиреля, Қаскелең, Қазақстан

ТОЧНОЕ И ПРИБЛИЖЕННОЕ РЕШЕНИЕ ДВУХФАЗОВОЙ ОБРАТНОЙ ЗАДАЧИ СТЕФАНА

Аннотация. Основная идея является нахождение точных решений двухфазной обратной задачи Стефана для вырожденной области с движущейся границей. Отслеживание ответов этих вопросов будут организованы следующим образом. Для нахождения аналитического решения мы в основном следуем методом, предложенным С.Н. Харин применяя формулу Фаа Ди Бруно для интегральных функций ошибок. В продолжении раздела интегральные функции ошибки и ее свойств, необходимых для разработки новых методов.

Задача Стефана очень сложна для точного решения. Поэтому мы пытаемся найти приближенное решение путем присвоения температурных профилей с изменением дифференциального уравнения на уравнения теплового баланса, которые получаются путем интегрирования по X и t . Для нахождения приближенного решения мы в основном следуем методом, предложенным С.Н. Харин, применения метода интегрального теплового баланса.

Ключевые слова: Метод интегральной функция ошибок, Метод интегрального теплового баланса, Двухфазовая обратная задача Стефана.

CONTENT

| | |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|
| <i>Baytanaev O.A.</i> Phenomenon of natural foci of zoonotic infections: a new hypothesis..... | 5 |
| <i>Turgumbayeva A.A., Rakhimov K.D., Ustenova G.O.</i> Antimicrobial and other medicinal properties of safflower (<i>carthamus tinctorius l.</i>) | 10 |
| <i>Abizhov M.M.</i> The beginning of cooperation between the countries of Russia and China..... | 15 |
| <i>Nursapa A.T.</i> Implementation of new economic policy: investment for bussinessmen | 21 |
| <i>Kudaikulov A.A., Josserand C., Kaltayev A.</i> Numerical investigation of fingering pattern formation during the flow of two immiscible fluids in a channel..... | 26 |
| <i>Rakhimbayeva A.A.</i> On the existing problems and perspectives of the eurasian economic union..... | 31 |
| <i>Sarsengeldin M., Kassabek S., Sagidolla B.</i> Exact and approximatesolutions of two phase inverse stefan problem..... | 37 |
| <i>Tulenbaev K.M., Shaimardanova Zh.N., Ghabdullin B.</i> Structure properties of (α, β) – bicommutative algebras..... | 49 |

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the work described has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the originality detection service Cross Check <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

www.nauka-nanrk.kz

<http://www.reports-science.kz/index.php/ru/>

Редакторы *М. С. Ахметова, Д. С. Аленов, Т.А. Апендиев*
Верстка на компьютере *А.М. Кульгинбаевой*

Подписано в печать 20.07.2016.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
11 п.л. Тираж 2000. Заказ 4.