

ISSN 2224-5227

2016 • 4

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
БАЯНДАМАЛАРЫ

ДОКЛАДЫ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН

REPORTS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN

ЖУРНАЛ 1944 ЖЫЛДАН ШЫҒА БАСТАҒАН

ЖУРНАЛ ИЗДАЕТСЯ С 1944 г.

PUBLISHED SINCE 1944



Бас редактор
ҚР ҰҒА академигі **М.Ж. Жұрынов**

Редакция алқасы:

хим.ғ. докторы, проф., ҚР ҰҒА академигі **Әдекенов С.М.** (бас редактордың орынбасары), эк.ғ. докторы, проф., ҚР ҰҒА академигі **Әділов Ж.М.**, мед. ғ. докторы, проф., ҚР ҰҒА академигі **Арзықұлов Ж.А.**, техн. ғ. докторы, проф., ҚР ҰҒА академигі **Бишімбаев У.К.**, а.-ш.ғ. докторы, проф., ҚР ҰҒА академигі **Есполов Т.И.**, техн. ғ. докторы, проф., ҚР ҰҒА академигі **Мұтанов Г.М.**, физ.-мат.ғ. докторы, проф., ҚР ҰҒА академигі **Өтелбаев М.О.**, пед. ғ. докторы, проф., ҚР ҰҒА академигі **Пралиев С.Ж.**, геогр.ғ. докторы, проф., ҚР ҰҒА академигі **Северский И.В.**; тарих.ғ. докторы, проф., ҚР ҰҒА академигі **Сыдықов Е.Б.**, физ.-мат.ғ. докторы, проф., ҚР ҰҒА академигі **Тәкібаев Н.Ж.**, физ.-мат.ғ. докторы, проф., ҚР ҰҒА академигі **Харин С.Н.**, тарих ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Әбүсейітова М.Х.**, экон. ғ. докторы, проф., ҰҒА корр. мүшесі **Бейсембетов И.К.**, биол. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Жамбакин К.Ж.**, тарих ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Кәрібаев Б.Б.**, мед. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Локшин В.Н.**, геол.-мин. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Өмірсеріков М.Ш.**, физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Рамазанов Т.С.**, физ.-мат. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Садыбеков М.А.**, хим.ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Сатаев М.И.**; ҚР ҰҒА құрметті мүшесі, а.-ш.ғ. докторы, проф. **Омбаев А.М.**

Редакция кеңесі:

Украинаның ҰҒА академигі **Гончарук В.В.** (Украина), Украинаның ҰҒА академигі **Неклюдов И.М.** (Украина), Беларусь Республикасының ҰҒА академигі **Гордиенко А.И.** (Беларусь), Молдова Республикасының ҰҒА академигі **Дука Г.** (Молдова), Тәжікстан Республикасының ҰҒА академигі **Илолов М.И.** (Тәжікстан), Қырғыз Республикасының ҰҒА академигі **Эркебаев А.Э.** (Қырғызстан), Ресей ҒА корр. мүшесі **Величкин В.И.** (Ресей Федерациясы); хим.ғ. докторы, профессор **Марек Сикорски** (Польша), тех.ғ. докторы, профессор **Потапов В.А.** (Украина), биол.ғ. докторы, профессор **Харун Парлар** (Германия), профессор **Гао Энджун** (КХР), филос. ғ. докторы, профессор **Стефано Перни** (Ұлыбритания), ғ. докторы, профессор **Богуслава Леска** (Польша), философия ғ. докторы, профессор **Полина Прокопович** (Ұлыбритания), профессор **Вуйцик Вольдемар** (Польша), профессор **Нур Изура Уздир** (Малайзия), д.х.н., профессор **Нараев В.Н.** (Ресей Федерациясы)

Главный редактор
академик НАН РК **М.Ж. Журинов**

Редакционная коллегия:

доктор хим. наук, проф., академик НАН РК **С.М. Адекенов** (заместитель главного редактора), доктор экон. наук, проф., академик НАН РК **Ж.М. Адилов**, доктор мед. наук, проф., академик НАН РК **Ж.А. Арзыкулов**, доктор техн. наук, проф., академик НАН РК **В.К. Бишимбаев**, доктор сельскохозяйств. наук, проф., академик НАН РК **Т.И. Есполов**, доктор техн. наук, проф., академик НАН РК **Г.М. Мутанов**, доктор физ.-мат. наук, проф., академик НАН РК **М.О. Отелбаев**, доктор пед. наук, проф., академик НАН РК **С.Ж. Пралиев**, доктор геогр. наук, проф., академик НАН РК **И.В. Северский**; доктор ист. наук, проф., академик НАН РК **Е.Б. Сыдыков**, доктор физ.-мат. наук, проф., академик НАН РК **Н.Ж. Такибаев**, доктор физ.-мат. наук, проф., академик НАН РК **С.Н. Харин**, доктор ист. наук, проф., чл.-корр. НАН РК **М.Х. Абусейтова**, доктор экон. наук, проф., чл.-корр. НАН РК **И.К. Бейсембетов**, доктор биол. наук, проф., чл.-корр. НАН РК **К.Ж. Жамбакин**, доктор ист. наук, проф., чл.-корр. НАН РК **Б.Б. Карибаев**, доктор мед. наук, проф., чл.-корр. НАН РК **В.Н. Локшин**, доктор геол.-мин. наук, проф., чл.-корр. НАН РК **М.Ш. Омирсериков**, доктор физ.-мат. наук, проф., чл.-корр. НАН РК **Т.С. Рамазанов**, доктор физ.-мат. наук, проф., чл.-корр. НАН РК **М.А. Садыбеков**, доктор хим. наук, проф., чл.-корр. НАН РК **М.И. Сатаев**; почетный член НАН РК, доктор сельскохозяйств. наук, проф., **А.М. Омбаев**

Редакционный совет:

академик НАН Украины **Гончарук В.В.** (Украина), академик НАН Украины **И.М. Неклюдов** (Украина), академик НАН Республики Беларусь **А.И. Гордиенко** (Беларусь), академик НАН Республики Молдова **Г. Дука** (Молдова), академик НАН Республики Таджикистан **М.И. Илолов** (Таджикистан), член-корреспондент РАН **Величкин В.И.** (Россия); академик НАН Кыргызской Республики **А.Э. Эркебаев** (Кыргызстан), д.х.н., профессор **Марек Сикорски** (Польша), д.т.н., профессор **В.А. Потапов** (Украина), д.б.н., профессор **Харун Парлар** (Германия), профессор **Гао Энджун** (КНР), доктор философии, профессор **Стефано Перни** (Великобритания), доктор наук, профессор **Богуслава Леска** (Польша), доктор философии, профессор **Полина Прокопович** (Великобритания), профессор **Вуйцик Вольдемар** (Польша), профессор **Нур Изура Уздир** (Малайзия), д.х.н., профессор **В.Н. Нараев** (Россия)

«Доклады Национальной академии наук Республики Казахстан» ISSN 2224-5227

Собственник: Республиканское общественное объединение «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №5540-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год. Тираж: 2000 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 218-220, тел. 272-13-19, 272-13-18

<http://nauka-nanrk.kz> reports-science.kz

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

©Национальная академия наук Республики Казахстан, 2016 г.

E d i t o r - i n - c h i e f

M.Zh. Zhurinov, academician of NAS RK

Editorial board:

S.M. Adekenov (deputy editor in chief), Doctor of Chemistry, prof., academician of NAS RK; **Zh.M. Adilov**, Doctor of Economics, prof., academician of NAS RK; **Zh.A. Arzykulov**, Doctor of Medicine, prof., academician of NAS RK; **V.K. Bishimbayev**, Doctor of Engineering, prof., academician of NAS RK; **T.I. Yespolov**, Doctor of Agriculture, prof., academician of NAS RK; **G.M. Mutanov**, Doctor of Physics and Mathematics, prof., academician of NAS RK; **M.O. Otelbayev**, Doctor of Physics and Mathematics, prof., academician of NAS RK; **S.Zh. Praliyev**, Doctor of Education, prof., academician of NAS RK; **I.V. Seversky**, Doctor of Geography, prof., academician of NAS RK; **Ye.B. Sydykov**, Doctor of Historical Sciences, prof., academician of NAS RK; **N.Zh. Takibayev**, Doctor of Physics and Mathematics, prof., academician of NAS RK; **S.N. Kharin**, Doctor of Physics and Mathematics, prof., academician of NAS RK; **M.Kh. Abuseitova**, Doctor of Historical Sciences, prof., corr. member of NAS RK; **I.K. Beisembetov**, Doctor of Economics, prof., corr. member of NAS RK; **K.Zh. Zhambakin**, Doctor of Biological Sciences, prof., corr. member of NAS RK; **B.B. Karibayev**, Doctor of Historical Sciences, prof., corr. member of NAS RK; **V.N. Lokshin**, Doctor of Medicine, prof., corr. member of NAS RK; **M.Sh. Omirserikov**, Doctor of Geology and Mineralogy, prof., corr. member of NAS RK; **T.S. Ramazanov**, Doctor of Physics and Mathematics, prof., corr. member of NAS RK; **M.A. Sadybekov**, Doctor of Physics and Mathematics, prof., corr. member of NAS RK; **M.I. Satayev**, Doctor of Chemistry, prof., corr. member of NAS RK; **A.M. Ombayev**, Honorary Member of NAS RK, Doctor of Agriculture, prof.

Editorial staff:

V.V. Goncharuk, NAS Ukraine academician (Ukraine); **I.M. Neklyudov**, NAS Ukraine academician (Ukraine); **A.I. Gordienko**, NAS RB academician (Belarus); **G. Duca**, NAS Moldova academician (Moldova); **M.I. Iolov**, NAS Tajikistan academician (Tajikistan); **A.E. Erkebayev**, NAS Kyrgyzstan academician (Kyrgyzstan); **V.I. Velichkin**, RAS corr.member (Russia); **Marek Sikorski**, Doctor of Chemistry, prof. (Poland); **V.A. Potapov**, Doctor of Engineering, prof. (Ukraine); **Harun Parlar**, Doctor of Biological Sciences, prof. (Germany); **Gao Endzhun**, prof. (PRC); **Stefano Perni**, Doctor of Philosophy, prof. (UK); **Boguslava Leska**, dr, prof. (Poland); **Pauline Prokopovich**, Doctor of Philosophy, prof. (UK); **Wójcik Waldemar**, prof. (Poland), **Nur Izura Udzir**, prof. (Malaysia), **V.N. Narayev**, Doctor of Chemistry, prof. (Russia)

Reports of the National Academy of Sciences of the Republic of Kazakhstan.

ISSN 2224-5227

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 5540-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 2000 copies

Editorial address: 28, Shevchenko str., of.219-220, Almaty, 050010, tel. 272-13-19, 272-13-18,

<http://nauka-nanrk.kz/> reports-science.kz

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

© National Academy of Sciences of the Republic of Kazakhstan, 2016

АНТИМИКРОБНЫЕ И ДРУГИЕ ЦЕЛЕБНЫЕ СВОЙСТВА САФЛОРЫ (*CARTHAMUS TINCTORIUS L.*)

А. А. Тургумбаева¹, К.Д. Рахимов², Г.О.Устенова¹

¹ Казахский Национальный медицинский университет им. С.Д. Асфендиярова, Алматы, Казахстан

² Казахский медицинский университет непрерывного образования, Алматы, Казахстан

e-mail: aknurik_88@mail.ru

Ключевые слова: *Carthamus tinctorius*, Asteraceae, сафлора, цветы, фитохимия, эфирное масло.

Впервые изучен состав цветков сафлоры и экспериментально подтверждена целесообразность использования полученных продуктов в составе лекарственных средств. Способ получения средства, обладающего антимикробной, противовоспалительной и ранозаживляющей средствами из цветки сафлоры (*Carthamus tinctorius L.*). Качественный состав эфирного масла цветки сафлоры и количественное содержание идентифицированных компонентов в нём устанавливали методом газо-жидкостной хроматографии GC-FID (Gas Chromatography – Flame Ionization Detector) с использованием стандартных образцов. Методом ГЖХ установлено наличие 8 компонентов, из них также идентифицированы по стандартным образцам beta-bisabolene 6,63 %, E-Nuciferol 9,06%, Z-Nuciferol 14,14 %, Cis-Lanceol 42, 45%.

УДК 615.322.012

ANTIMICROBIAL AND OTHER MEDICINAL PROPERTIES OF SAFFLOWER (*CARTHAMUS TINCTORIUS L.*)

A.A.Turgumbayeva¹, K.D. Rakhimov², G.O. Ustenova¹

¹S.D. Asfendiyarov Kazakh National Medical University, Almaty, Republic of Kazakhstan

²Kazakh Medical University Continuing Education, Almaty, Kazakhstan

Keywords: *Carthamus tinctorius*, Asteraceae, safflower, flowers, phytochemistry, volatile oil.

Abstract: *Carthamus tinctorius L. (Asteraceae)* is used medicinally in Europe, China and India amongst several places in the world. The aim was designed to study the biological activity and chemical composition of volatile oil of *Carthamus tinctorius L.* The composition of the volatile oil obtained from the dried flowers of *Carthamus tinctorius L.* collected in Kazakhstan was analyzed by gas chromatography (GC) and gas chromatography-mass spectrophotometry (GC- FID). 8 known and 2 unknown compounds were detected from the extract. The major compounds of the oil were 3-carene, beta-bisabolene, alpha-trans-bergamotol, Z-nuciferol, E-nuciferol, cis-Lanceol, n-tricosane and pentacosane. The volatile oil of this plant have been demonstrated to possess multiple pharmacological activities. In this review, we have explored the phytochemistry and pharmacological activities of *CarthamusTinctorius L.* as well as have received antimicrobial, anti-inflammatory, a medicinal ointment.

Introduction. Safflower, *Carthamus tinctorius L.*, is a member of the family Compositae or Asteraceae, cultivated mainly for its seed, which is used as edible oil and as birdseed. Traditionally, the crop was grown for its flowers, used for colouring and flavouring foods and making dyes, especially before cheaper aniline dyes became available, and in medicines.

The plant has a strong taproot which enables it to thrive in dry climates. Safflower is one of humanity's oldest crops, but generally it has been grown on small plots for the grower's personal use and

it remains a minor crop with world seed production around 800 000 t per year. Oil has been produced commercially and for export for about 50 years, first as an oil source for the paint industry, now for its edible oil for cooking, margarine and salad oil. Over 60 countries grow safflower, but over half is produced in India (mainly for the domestic vegetable oil market). Production in the USA, Mexico, Ethiopia, Argentina and Australia comprises most of the remainder. China has a significant area planted to safflower, but the florets are harvested for use in traditional medicines and the crop is not reported internationally. Also, the crop has also been cultivated in many other countries, such as Kazakhstan, Ethiopia, Argentina, China, Uzbekistan, Australia, Russian Federation, Pakistan, and Spain [1, 2]. Traditionally, the crop was grown for its flowers, used for coloring and flavoring foods and making dyes, especially before cheaper aniline dyes became available and in medicines. It is considered one of the alternative oil crops, particularly in the dry and semi dry lands due to its tolerance to drought, salinity and cold stress. Safflower oil quality is high due to its fatty acids composition. Standard safflower oil contains about 6-8 % palmitic acid, 2-3 % stearic acid, 16-20% oleic acid and 71-75 % linoleic acid. In addition, very low levels of myristic (0.24 %) and behenic (0.43 %) acids were recorded in its oil [3].

Safflower is widely distributed in eastern and western Asia. The flower of Safflower is used in folk medicine as an analgesic, antithrombotic and antihypertensive crude drug as well as a source of natural colorants [4-8]. Safflower has long been grown for the dye extracted from the flowers. Depending on the dyeing procedure and the addition of other colorants and mordant's, it imparts a yellow, red, brown or purple color to cloth. With the introduction of cheap synthetic dyes, its importance as a dye source has greatly declined. However, dyes are still produced on a small scale for traditional and religious purposes.

Materials and methods

Plant Material: The plant materials used in study were obtained from Almaty, southern Kazakhstan, safflower (*Carthamus tinctorius L.*) collected in the summer. The plant was identified by taxonomist Konyrbekov M. of the station. A voucher specimen was preserved at the herbarium Krasnovodopadskaya Breeding Experimental Station, Ministry of Agriculture, Republic of Kazakhstan.

Phytochemistry

More than 200 compounds have been isolated from *C. tinctorius* and the commonly known ones are flavonoids, phenylethanoid glycosides, coumarins, fatty acids, steroids and polysaccharides [9]. Analysis of safflower seeds showed that crude protein ranged from 14.9 % to 17 %, total sugar from 3.2 % to 9.2 % and extractable lipids from 25 % to 40 % [10]. Oil content of the seeds is similar to that of olive and includes linoleic acid (63 %–72 %), oleic acid (16 %–25 %), and linolenic acid (1 %–6 %) [11-12].

Results and discussion

Compounds were quantified by performing area percentage calculations based on the total combined FID area. For example, the area for each reported peak was divided by total integrated area from the FID chromatogram from all reported peaks and multiplied by 100 to arrive at a percentage. The percentage of a peak is a percentage relative to all other constituents integrated in the FID chromatogram. The 8 components represented about 98.642% of the total detected constituents. The percentage content of the individual components, retention time, Kovat's index (KI) are summarized in Table 1. The differences in chemical composition essential oil of the present study and previous research may be because of the geographic and climatic factors, chemo types, drying conditions and mode of distillation.

Table 1.0: Chemical composition of *Carthamus tinctorius L*

Compounds	Retention time	Kovat Index (KI)	Area %
3-Carene	8.325	1007.70	0.967
Beta-Bisabolene	28.546	1512.75	6.629
Unknown	29.113	1526.91	1.392
Alpha-trans-bergamotol	35.195	1686.46	3.457
Z-Nuciferol	36.585	1725.99	14.139
E-Nuciferol	37.713	1755.92	9.057
Cis-Lanceol	37.896	1761.72	42.449
Unknown	44.465	1957.40	11.967
n-Tricosane	54.610	2299.22	4.197
Pentacosane	59.914		4.388

Antimicrobial activity. Antimicrobial study was done for the volatile oil against five fungi and five bacteria. The oil was found to have a good activity against *Cryptococcus neoformans* ATCC 90113 with an IC₅₀ value of 8 µg/ml. However, no antimicrobial activity was shown against *Carthamus tinctorius* L (Table 2). [6, 13], reported that the essential oil of *Carthamustinctorius*L exhibited a negative antifungal effect using broth microdilution and disc gel diffusion methods. The antifungal activity was assessed against five dermatophytes (*Trichophyton mentagrophytes*, *T.rubrum*, *Microsporumcanis*, *M.nanum* and *Epidermophytonfloccosum*), three filamentous fungi (*Aspergillousniger*, *A.fumigatus* and *Mucorsps.*) and five strains of yeast (*Saccharomyces cerevisiae*, *C.neoformans*, *Candida albicans*, *C.tropicalis* and *Torulopsisglabrata*). This report is similar with what we are reporting however we found the volatile oil *Carthamustinctorius*L IC₅₀ value of 20.41 µg/ml against *C.neoformans*.

Antimalarial activity. Antimalarial activity was studied for the volatile oil of *Carthamustinctorius*L against chloroquine sensitive *Plasmodium falciparum* (D6, Sierra Leone) and resistant (W2, Indo China). The oil showed moderate antimalarial activity (IC₅₀ 47600 µg/ml against *P. falciparum* D6 and *P. falciparum* W2). *Carthamustinctorius*L showed good activity with IC₅₀ values >47600 µg/ml against *P. falciparum* D6 and *P. falciparum* W2 (Table 2) [14].

Antileishmanial activity. Antileishmanial evaluation was done on the oil *Carthamustinctorius*L against *Leishmania donovani*. The oil showed moderate activity with IC₅₀ and IC₉₀ values of 80.0µg/ml. *Carthamustinctorius*L showed activity with IC₅₀ and IC₉₀>80 [15].

Table 2.0 – Biological activity of essential oil of *Carthamus tinctorius* L.

Biological activity	Test parasite	<i>Carthamustinctorius</i> L essential oil (µg/ml)
Antileishmanial activity	<i>L. donovani</i>	
	IC ₅₀	>80
	IC ₉₀	>80
Antimalarial activity	<i>P.falciparum</i> D6	
	IC ₅₀	>47600
	W2 IC ₅₀	
Antimicrobial	<i>C. neoformans</i>	<20.41

An ointment of essential oil the from safflower flowers grown Kazakhstan: for creating ointment of flowers safflower (*Carthamus tinctorius* L.) optimum composition of the excipients. So several models were created ointment bases - emulsion, a slurry, combined with application of various the excipients - sunflower oil, glycerol, paraffin oil, lanolin, etc., Emulsifiers - Tween-80, T-2 and others. The most efficient composition of the technological parameters was ointment base with the following composition (Table 3):

Table 3.0 The composition an ointment consisting 100 g.

The active substance	
essential oil obtained from the flowers of safflower	9.0
The auxiliary substances	
sunflower oil	40.0
T-2	5.0
Purified Water	46.8
<i>Oleum Menthaepiperitae</i>	0.2
Total weight	100, 0

By its consistency suitable for application of ointment and a prolonged exposure to lesion focus. Other compositions were not suitable for their consistency of, as were liquefaction or thick, which was inconvenient terms of application, and distinguished by the fact that the system was subjected to separation and was losing their structural and mechanical properties.

At the stage of carried out preparation and auxiliary of drugs and materials. At the initial stage emulsion base was prepared of the following composition: 40 parts of sunflower oil, 5 parts of emulsifier T-2 ad 100.0 parts of purified water. Emulsion base was prepared in the following way: an emulsifier T-2 was melted in a water bath, and added to overheated oil in the last turn slowly added in a thin stream hot water (90 °C). Then, medicinal substances were injected into the finished emulsion base. The addition of

essential oil obtained from safflower flowers to the base was carried out at preliminary maximum dispersing them before molecular state, which was carried out with constant stirring to forming a stable system. In the next step was carried out homogenization ointment to obtain homogeneous mass. Ointment from almost white to slightly yellowish color with a weak a characteristic smell of *Oleum Menthaepiperitae*. The proposed method allows to obtain means of wound healing, regenerating and anti-inflammatory activity.

The indexes of quality according with the requirements of the State Pharmacopoeia of the Republic of Kazakhstan.

Conclusions

From the result of the study, it could be concluded that the safflower collected from the Southern region of Kazakhstan is one of the best genotype available. *Carthamus tinctorius* is regarded as a valuable plant in Kazakh system of medicine, Chinese medicine and modern drug development areas for its versatile medicinal uses. The aim was designed to study the biological activity and chemical composition of volatile oil of *Carthamus tinctorius L.* The composition of the volatile oil obtained from the dried flowers of *Carthamus tinctorius L.* growing in Kazakhstan was analyzed by gas chromatography (GC) and gas chromatography-mass spectrophotometry (GC- FID). 8 known and 2 unknown compounds were detected from the extract. The major compounds of the oil were 3-carene, beta-bisabolene, alpha-transbergamotol, Z-nuciferol, E-nuciferol, cis-Lanceol, n-tricosane and pentacosane. The traditional use of *Carthamus tinctorius L.* against various skin infections wounds has been corroborated since the extracts displayed in vitro antimicrobial properties against different test organisms. The fact that the ethyl acetate extracts of this medicinal plant were very active against the test organisms. To our knowledge from literature, this is the first time, report on the antileishmanial activity, antimalarial activity and antimicrobial properties of extracts. Obtained an experimental industrial series of ointment based medicinal vegetative raw materials (*Carthamus tinctorius L.*).

Development of an optimal composition and rational technology medicinal products based on essential oil obtained from the flowers of safflower, sunflower oil, emulsifiers T-2 and others. The antimicrobial, anti-inflammatory, regenerative, curative effect. From the result obtained in the study traditional use of *Carthamus tinctorius L.* against various skin infections wounds has been corroborated since the extracts displayed in vitro antimicrobial properties against different test organisms. Further studies of other phyto-active compounds will possibly lead to exploration of new methods for therapeutic and industrial application.

REFERENCES

- [1] Dajue L. and Mundel H. H., Safflower (*Carthamus tinctoriusL.*). Promoting the conservation and use of underutilized and neglected crops, Monogr // 7. Institut für Pflanzengenetik and Kultur. (IPK), Gatersleben, Germany, and International Plant Genetic Resources Institute, Rome, Italy, 8. -1996. –P. 32-39.
- [2] Turgumbayeva A.A., Ustenova G.O., Samir A Ross. Volatile Oil Composition of *Carthamus Tinctorius L* the Flowers Grown in Kazakhstan // Research Journal of Pharmaceutical, Biological and Chemical Sciences. -2015. - Vol. 6. –Iss. 2. – P. 125-129.
- [3] Belgin C., Bilal G., Mustafa K. Inter. J. Nat. and Eng. Sci., - 2007. 1(3): 11-15.
- [4] Zhou F.R., Zhao M.B., Tu P.F. Simultaneous determination of four nucleosides in *Carthamus tinctorius L.* and Safflower injection using high performance liquid chromatography // J Chin Pharmaceut. Sci (Chin). - 2009. -P. 326-330.
- [5] Kruawan K., Kangsadalampai K., Antioxidant activity, phenolic compound contents and antimutagenic activity of some water extract of herbs // Thai J. Pharm. Sci. -2006. 30:28-35.
- [6] Jun M.S., Ha Y.M., Kim H.S., Jang H.J., Kim Y.M., Lee Y.S., et al. Anti-inflammatory action of methanol extract of *Carthamus tinctorius* involves in heme oxygenase-1 induction // J. Ethnopharmacol. - 2011. 133:524-530.
- [7] Almeida R.N., Navarro D.S., Barbosa-Filho J.M. Plants with central analgesic activity // Phytomed. – 2001. 8:310-322.
- [8] Kasahara Y., Kumaki K., Sato T., Katagiri S. Pharmacological studies on flower petals of *Carthamus tinctorius* central actions and anti-inflammation. -1989. 43:331-338.
- [9] Zhou F.R., Zhao M.B., Tu P.F. Simultaneous determination of four nucleosides in *Carthamus tinctorius L.* and Safflower injection using high performance liquid chromatography // J. Chin. Pharmaceut. Sci (Chin). – 2009. 18:326-30.
- [10] Kim S.K., Cha J.Y., Jeong S.J., Chung C.H., Choi Y.R., Cho Y.S. Properties of the chemical composition of safflower (*Carthamus tinctorius L.*) sprout // Korean J. Life Sci. – 2000. 10:68-73.
- [11] Lee J.Y., Chang E.J., Kim H.J., Park J.H., Choi S.W. Antioxidative flavonoids from leaves of *Carthamus tinctorius*. Arch. Pharm. Res. – 2002. 25:313-19.
- [12] Bagheri H., Andalibi B., Zangani E., Jamshid S. Safflower (*Carthamus tinctorius* cv. Sina) oil and seed yield

improvement in rainfed condition by atrazine foliar application // Ann.Biol.Res. – 2012. 3(2):1202-9.

[13] Ma G., Khan S.I., Jacob M.R., Tekwani B.L., Li Z., Pasco D.S., Walker L.A., I.A. Khan, Antimicrobials and antileishmanial activities of hypocrellins A and B. Antimicrob // Agents Chemother. – 2004. 48:445-450.

[14] Makler M.T., and Hinrichs D.J., Measurement of lactate dehydrogenase activity of Plasmodium falciparum as an assessment of parasitemia // The Am. J. Trop. Med. Hyg. – 1993. 48: 205-210.

[15] Okada Y., and Okada M., Scavenging effect of soluble proteins in broad beans on free radicals and active oxygen species // J. Agric. Food Chem., -1998. 46: 401-406.

МАҚСАРЫ ӨСІМДІГІНІҢ (*CARTHAMUS TINCTORIUS L.*) МИКРОБҚА ҚАРСЫ ЖӘНЕ БАСҚА ШИПАЛЫ ҚАСИЕТТЕРІН ЗЕРТТЕУ

А. А. Тургумбаева¹, К.Д. Рахимов², Г.О.Устенова¹.

¹ С.Ж.Асфендияров атындағы Қазақ Ұлттық Медицина Университеті
Алматы, Қазақстан.

² Қазақ медициналық үздіксіз білім беру университет, Алматы, Қазақстан.

Тірек сөздер: *Carthamus tinctorius*, Asteraceae, мақсары, гүл, фитохимия, эфир майы.

Алғаш рет мақсары өсімдігінің гүлі тәжірбие жүзінде зерттелді және дәрілік қалып жасау негіздемесі ұсынылды. Шикізат гүлінен *Carthamus Tinctorius* микробқа, қабынуға қарсы және жара жазушы әсері бар дәрілік қалып жасады. Сол себепті біздің зерттеуіміздегі Қазақстандық мақсары өсімдігінің гүлінен сары түсті, майлы экстракт алынды. Өсімдік шикізаты гүлінен алынған майлы экстрактының эфир майлары құрамы GC-FID (Gas Chromatography – Flame Ionization Detector, газ хроматограммасы жалынды фотометорлық детектор) анықталды, нәтижесінде келесідей заттар көп бөлінді beta-bisabolene 6,63 % , E-Nuciferol 9,06%, Z-Nuciferol 14,14 % , Cis-Lanceol 42, 45%.

Поступила 26.06.2013 г.

СОДЕРЖАНИЕ

Математика	
<i>Мартынов Н.И., Рамазанова М.А.</i> Краевые задачи обобщенной плоской деформации линейно - упругого тела.....	5
Технические науки	
<i>Алтынбек Ш.Ч., Байконурова А.О.</i> Изучение влияния состава золотосодержащих растворов на сорбцию золота природными и синтетическими ионитами.....	17
Химия	
<i>Гылымхан Н.Т., Жумагалиева Ш.Н., Абилов Ж.А.</i> Возможности использования бентонитовых глин в медицине.....	24
<i>Мальшиев В.П., Зубрина Ю.С., Макашева А.М., Каткеева Г.Л., Кайкенов Д.А.</i> Особая роль и более общее выражение стерического фактора в вероятностной модели измельчения.....	34
<i>Хусанов А.Е., Крамалев С.О., Калдыбаева Б.М., Абилямагжанов А.З.</i> Исследование эмульгирующих свойств продуктов амидирования растительных масел.....	45
Науки о Земле	
<i>Абаканов Т., Садыкова А.Б., Ли А.Н., Степаненко Н.П.</i> Формализованная оценка сейсмopotенциала (M_{max}) земной коры Казахстана по комплексу сейсмогеофизических параметров.....	53
<i>Исламкулов К.М., Мырхалыков Ж.У.</i> Повышение прочности и износостойкости металлорежущих инструментов и литых деталей.....	61
Биология	
<i>Мукушкина Д.Д., Шертай М.Ж., Мирошник Т.Н., Аширбеков Е.Е., Талаева Ш.Ж., Хансеитова А.К., Айтхожина Н.А.</i> Выявление ассоциации полиморфизмов гена <i>ESR1</i> с риском развития рмж среди населения Казахстана.....	67
<i>Байтанаев О. А.</i> Феномен природной очаговости зоонозных инфекций: новая гипотеза.....	74
<i>Рсалиев А.С., Чудинов В.А., Амирханова Н.Т.</i> Устойчивость селекционных материалов ячменя Карабалыкской сельскохозяйственной опытной станции к сетчатой пятнистости и мучнистой росе.....	79
Медицина	
<i>Тургумбаева А. А., Рахимов К.Д., Устенова Г.О.</i> Антимикробные и другие целебные свойства сафлоры (<i>CARTHAMUS TINCTORIUS L.</i>).....	88
<i>Качиева З.С., Габдулина Г.Х.</i> Роль полиморфизмов гена <i>dkk1</i> в развитии деградации сустава у пациентов с ревматоидным артритом.....	93
Аграрные науки	
<i>Кулажанов Т.К.</i> Определение выхода мяса при национальной разделке туши баранины.....	100
Общественные науки	
<i>Абижов М. М.</i> Начало сотрудничества стран России и Китая.....	105
<i>Аюпова З.К., Кусаинов Д.У.</i> К вопросу об особенностях образования в мировой культуре.....	110
<i>Довгань А. В.</i> Смысл как детерминанта религии (на примере христианства).....	119
<i>Кишибекова Г.К., Абдулина Г.А., Жанбырбаева С.М.</i> Факторы повышения конкурентоспособности национальной экономики в условиях глобализации.....	124
<i>Кулбай Б.С.</i> Разработка адаптированной модели управления предприятием текстильной промышленности.....	135
<i>Курманов Н.А., Бактымбет С.С., Бактымбет А.С.</i> Развитие человеческого потенциала в Казахстане и в странах евразийского экономического союза.....	141
<i>Смаилова Ж.П., Таспенова Г.А., Карымсакова Ж.К.</i> Свободно плавающий валютный курс как инструмент денежно-кредитной политики.....	148
<i>Раимбердиев Т.П., Нысанбаева А.М., Носаненко Г.</i> Роль менеджмента в реформировании институтов государственного управления.....	154
<i>Рахимбекова А.Е., Курманов Н.А., Махатова А.Б., Серикбаева Э.А.</i> Развитие фармацевтической отрасли в республике Казахстан как основного рыночного элемента системы здравоохранения.....	161
<i>Татибеков Б. Л., Аман Р.Л.</i> Основные принципы сокращения неформального рынка труда и неформальной занятости на современном этапе.....	171