

ISSN 2518-1483 (Online),
ISSN 2224-5227 (Print)

2017 • 1

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

БАЯНДАМАЛАРЫ

ДОКЛАДЫ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН

REPORTS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN

ЖУРНАЛ 1944 ЖЫЛДАН ШЫҒА БАСТАҒАН

ЖУРНАЛ ИЗДАЕТСЯ С 1944 г.

PUBLISHED SINCE 1944



Б а с р е д а к т о р ы
х.ғ.д., проф., ҚР ҰҒА академигі **М.Ж. Жұрынов**

Р е д а к ц и я а л қ а с ы:

Адекенов С.М. проф., академик (Қазақстан) (бас ред. орынбасары)
Боос Э.Г. проф., академик (Қазақстан)
Величкин В.И. проф., корр.-мүшесі (Ресей)
Вольдемар Вуйчик проф. (Польша)
Гончарук В.В. проф., академик (Украина)
Гордиенко А.И. проф., академик (Белорус)
Дука Г. проф., академик (Молдова)
Илолов М.И. проф., академик (Тәжікстан),
Леска Богуслава проф. (Польша),
Локшин В.Н. проф. чл.-корр. (Қазақстан)
Нараев В.Н. проф. (Ресей)
Неклюдов И.М. проф., академик (Украина)
Нур Изура Удзир проф. (Малайзия)
Перни Стефано проф. (Ұлыбритания)
Потапов В.А. проф. (Украина)
Прокопович Полина проф. (Ұлыбритания)
Омбаев А.М. проф. (Қазақстан)
Өтелбаев М.О. проф., академик (Қазақстан)
Садыбеков М.А. проф., корр.-мүшесі (Қазақстан)
Сатаев М.И. проф., корр.-мүшесі (Қазақстан)
Северский И.В. проф., академик (Қазақстан)
Сикорски Марек проф. (Польша)
Рамазанов Т.С. проф., корр.-мүшесі (Қазақстан)
Такибаев Н.Ж. проф., академик (Қазақстан), бас ред. орынбасары
Харин С.Н. проф., академик (Қазақстан)
Чечин Л.М. проф., корр.-мүшесі (Қазақстан)
Харун Парлар проф. (Германия)
Энджун Гао проф. (Қытай)
Эркебаев А.Э. проф., академик (Қырғыстан)

«Қазақстан Республикасы Ұлттық ғылым академиясының баяндамалары»
ISSN 2518-1483 (Online),
ISSN 2224-5227 (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» Республикалық қоғамдық бірлестігі (Алматы қ.)
Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде 01.06.2006 ж.
берілген №5540-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.
Тиражы: 2000 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
<http://nauka-nanrk.kz>, reports-science.kz

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2017

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Главный редактор
д.х.н., проф., академик НАН РК **М. Ж. Журинов**

Редакционная коллегия:

Адекенов С.М. проф., академик (Казахстан) (зам. гл. ред.)
Боос Э.Г. проф., академик (Казахстан)
Величкин В.И. проф., чл.-корр. (Россия)
Вольдемар Вуйчик проф. (Польша)
Гончарук В.В. проф., академик (Украина)
Гордиенко А.И. проф., академик (Беларусь)
Дука Г. проф., академик (Молдова)
Илолов М.И. проф., академик (Таджикистан),
Леска Богуслава проф. (Польша),
Локшин В.Н. проф. чл.-корр. (Казахстан)
Нараев В.Н. проф. (Россия)
Неклюдов И.М. проф., академик (Украина)
Нур Изура Удзир проф. (Малайзия)
Перни Стефано проф. (Великобритания)
Потапов В.А. проф. (Украина)
Прокопович Полина проф. (Великобритания)
Омбаев А.М. проф. (Казахстан)
Отелбаев М.О. проф., академик (Казахстан)
Садыбеков М.А. проф., чл.-корр. (Казахстан)
Сатаев М.И. проф., чл.-корр. (Казахстан)
Северский И.В. проф., академик (Казахстан)
Сикорски Марек проф., (Польша)
Рамазанов Т.С. проф., чл.-корр. (Казахстан)
Такибаев Н.Ж. проф., академик (Казахстан), зам. гл. ред.
Харин С.Н. проф., академик (Казахстан)
Чечин Л.М. проф., чл.-корр. (Казахстан)
Харун Парлар проф. (Германия)
Энджун Гао проф. (Китай)
Эркебаев А.Э. проф., академик (Кыргызстан)

Доклады Национальной академии наук Республики Казахстан»

ISSN 2518-1483 (Online),

ISSN 2224-5227 (Print)

Собственник: Республиканское общественное объединение «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №5540-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год.

Тираж: 2000 экземпляров

Адрес редакции: 050010, г.Алматы, ул.Шевченко, 28, ком.218-220, тел. 272-13-19, 272-13-18

<http://nauka-nanrk.kz> reports-science.kz

©Национальная академия наук Республики Казахстан, 2017 г.

Адрес типографии: ИП «Аруна», г.Алматы, ул.Муратбаева, 75

E d i t o r i n c h i e fdoctor of chemistry, professor, academician of NAS RK **M.Zh. Zhurinov****E d i t o r i a l b o a r d :****Adekenov S.M.** prof., academician (Kazakhstan) (deputy editor in chief)**Boos E.G.** prof., academician (Kazakhstan)**Velichkin V.I.** prof., corr. member (Russia)**Voitsik Valdemar** prof. (Poland)**Goncharuk V.V.** prof., academician (Ukraine)**Gordiyenko A.I.** prof., academician (Belarus)**Duka G.** prof., academician (Moldova)**Ilov M.I.** prof., academician (Tadjikistan),**Leska Boguslava** prof. (Poland),**Lokshin V.N.** prof., corr. member. (Kazakhstan)**Narayev V.N.** prof. (Russia)**Nekludov I.M.** prof., academician (Ukraine)**Nur Izura Udzir** prof. (Malaysia)**Perni Stephano** prof. (Great Britain)**Potapov V.A.** prof. (Ukraine)**Prokopovich Polina** prof. (Great Britain)**Ombayev A.M.** prof. (Kazakhstan)**Otelbayv M.O.** prof., academician (Kazakhstan)**Sadybekov M.A.** prof., corr. member. (Kazakhstan)**Satayev M.I.** prof., corr. member. (Kazakhstan)**Severskiy I.V.** prof., academician (Kazakhstan)**Sikorski Marek** prof., (Poland)**Ramazanov T.S.** prof., corr. member. (Kazakhstan)**Takibayev N.Zh.** prof., academician (Kazakhstan), deputy editor in chief**Kharin S.N.** prof., academician (Kazakhstan)**Chechin L.M.** prof., corr. member. (Kazakhstan)**Kharun Parlar** prof. (Germany)**Endzhun Gao** prof. (China)**Erkebayev A.Ye.** prof., academician (Kyrgyzstan)**Reports of the National Academy of Sciences of the Republic of Kazakhstan.****ISSN 2224-5227****ISSN 2518-1483 (Online),****ISSN 2224-5227 (Print)**

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 5540-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 2000 copies

Editorial address: 28, Shevchenko str., of.219-220, Almaty, 050010, tel. 272-13-19, 272-13-18,
<http://nauka-nanrk.kz / reports-science.kz>

© National Academy of Sciences of the Republic of Kazakhstan, 2017

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

**REPORTS OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN**

ISSN 2224-5227

Volume 1, Number 311 (2017), 69 – 77

UDC 502.53/57.087

V. Sadomskiy¹, E. Krupa², I. Aminova³¹“SED” LLP, Almaty, Kazakhstan;²Republican State Enterprise “Institute of Zoology”, Almaty, Kazakhstan;³“Akjayik” State Nature Reserve, Atyrau, Kazakhstane-mail: sadomsky@list.ru, elena_krupa@mail.ru, agrimony92@mail.ru**EXPERIMENTAL RESEARCH OF SEISMIC-ACOUSTIC
IMPACT ON THE NORTH CASPIAN AQUATIC ORGANISMS**

Abstract. Materials of experimental and field surveys of seismic-acoustic impact on the North Caspian zooplankton, zoobenthos and fish fauna in summer, 2012-2013 are provided in this article. The article also demonstrates that for assessment of seismic exploration impact, the following is of high indicator value: proportion of traumatized or dead specimens, the values of Shannon-Weaver diversity index and the value of an average individual mass of the specimen in hydrocoenosis. The highest proportion of the traumatized and/or dead specimens was recorded in plankton and benthic invertebrates' populations in the course of experimental tests at a distance of 1m and 5m from the seismic source. At the same period, a deviation of the Shannon-Weaver diversity index values and average individual mass of the specimen from background values was recorded in the both communities. Species composition of fish fauna was characterized by a high level of similarity throughout all stages of the surveys implemented. The lowest indicators of diversity (number of species, values of Shannon-Weaver index) and fish numbers were recorded during seismic exploration. An increased average individual mass of a specimen in ichthyocoenosis together with the reduced numbers during seismic exploration may indicate the avoidance of unfavorable zone by younger fishes.

Keywords: zooplankton, zoobenthos, fishes, seismic acoustics, pneumatic source, seismic surveys.

An intense seismic exploration aimed at development of oil fields is causing the increased anthropogenic load on the entire Caspian Sea ecosystem [1]. Currently published data on assessment of seismic-acoustic impact do not present the level and scope of its effect on various groups of aquatic organisms constituting the relevance of this article [2-8].

Experimental study of seismic-acoustic effects on aquatic communities was implemented in two North Caspian areas (Figure 1) in June, 2012 and August, 2013. Three studies – before impact (background level, stage 1), during pneumatic source impact (experimental tests, stage 2) and after completion of seismic exploration impact (stage 3). To evaluate the impact of seismic acoustic effects on behavioral response of fish fauna, the composition and quantity of bento-pelagic fish communities were studied throughout all stages of seismic surveys.

Selection and processing of zooplankton, zoobenthos and fish fauna samples were performed using standard methods [1-5]. At all stages of the surveys, proportion of injured and/or dead specimens was studied in the populations of plankton and benthic invertebrates.

During experimental tests, cages with the fish and zooplankton samples caught in advance were placed at the depth of 3-4 m and at a distance of 1 m, 5 m and 10 m from the pneumatic source (Figure 2). In order to maintain the relative accuracy in the distance between the pneumatic source and cages setting points, a seismic-acoustic excitation signal was performed by the pontoon device towed by shooter at the time of its passage through the reference buoy. Sediment samples with zoobenthos were collected on reference stations within 15 minutes after the pneumatic source seismic acoustic signals.

Hydrological and hydro-physical parameters were registered simultaneously with the experimental tests.




Figure 1. Layout of test sites according to the assessment of the impact seismic survey on the North Caspian aquatic organisms



Figure 2 - Scheme of the location of stations to study the impact of seismoacoustics on aquatic organisms

The experimental areas were characterized by shallow depths with similar hydrophysical and hydrochemical characteristics (Table 1). The sediments were presented by terrigenous sediments with sand fraction dominated (76.5-54.6%) (0.05-1.0 mm) [6].

Table 1 – Hydrological and hydrophysical indicators on experimental areas

Areas	Depth, m	Transparency, m	Temperature, °C	Water salinity, ‰	Turbidity, NTU	pH	Flow rate, m/sec
Area No.1, June 2012	6.5	1.0	26.6	5.1	9.06	7.7	9-11
Area No.2, August 2013	5.1	0.7	24.8	6.0	8.9	8.4	7-13

The diversity of zooplankton was ranged from 23 to 50 taxa. The background was typical of the Caspian Sea species: rotifers *Brachionus quadridentatus*, *B. plicatilis*, *Filinia longiseta*; cladocerans *Cornigerius maeoticus hirsus*, *Podonevadne camptonyx*, *P. angusta*, *P. trigona*, *Cercopagis pengoi*; copepods *Acartia tonsa*, *Calanipeda aquedulcis*: larvae *Cirripedia*, *Bivalvia*, *Hediste diversicolor*; crab *Rhitropanopeus harrisi*. The composition of the background species preserved substantial similarity throughout all stages of the surveys.

The number of species of planktonic invertebrates and their total number in the background conditions and after impact were characterized by the same or similar values (Table 2). During the experimental tests, a low diversity and deviation of zooplankton abundance from the values obtained in stages 1 and 3 was according to the survey methodology – staying of experimental communities in isolated cages.

Relative indicators have most indicator significance – the proportion of injured or dead specimens, the values of diversity index of Shannon-Weaver and average individual weight of specimen. The proportion of injured or dead specimens in the plankton populations of invertebrates at the background level, and after the impact was approximately the same, or lowers than at the time of impact seismoacoustics. The incidence of dead and/or injured specimens of copepod *Calanipeda aquedulcis* and *Acartia tonsa* at all stages of the surveys was close in value to the other species, except for *H. diversicolor* polychaete and *B. plicatilis* rotifers, incidence of dead and/or injured specimens was higher during impact.

Table 2 – Structural indicators of the North Caspian zooplankton at various survey stages

Indicator	Background conditions	Experimental tests	After impact
¹ number of species	34	29	35
² number of species	19	10	18
¹ average abundance, thousand spec./m ³	43.5	110.0	53.9
² average abundance, thousand spec./m ³	95.6	31.0	71.7
¹ proportion of injured or dead specimens, %	0.5	10.8	0.7
¹ Shannon-Weaver index, bit/ spec.	2.12	1.84	2.14
² Shannon-Weaver index, bit/ spec.	1.94	2.36	1.86
¹ Shannon-Weaver index, bit/mg	1.88	2.32	2.05
² Shannon-Weaver index, bit/mg	0.74	0.44	1.27
¹ average individual masse, mg	0.0051	0.0045	0.0051
² average individual masse, mg	0.0540	0.0100	0.0470
Note: numerals indicate the corresponding areas			

Analysis of the spatial distribution showed that the proportion of injured and/or dead planktonic invertebrate species was highest at a distance from the pneumatic source from 1 to 5 m. At a distance of 10 meters, it decreased with varying intensity, but did not reach background values (Figure 3).

The dynamics of the Shannon-Weaver index at each area had its own characteristics (Figure 4). The values of the indicator, according to the number of types of proportion in the total population (bits/sp.), in the background period and after the impact of seismic operations on both areas did not differ. Thus, during the experimental tests, the index value recorded in the area No.1 has decreased, and in the area No.2, in contrast, has increased with respect to the other two stages.




Figure 3 – Changes in proportion of injured and/or dead specimens in zooplankton at different distance from the pneumatic source

During the experimental tests, the dynamics of the second option of the index (bits/mg) was characterized by opposing trends: in the area No.1, it was positive, and in the area No.2, it was negative. At the same time, the indicator of relative background level was slightly higher than it was recorded after the impact completion.




Figure 4 – The dynamics of the Shannon-Weaver index values at different stages of surveys of seismoacoustics impact on zooplankton

Table 3 – Structural indicators of the North Caspian zoobenthos at various survey stages

Indicator	Background conditions	Experimental tests	After impact
¹ number of species	20	14	13
² number of species	14	15	10
¹ average abundance, thousand spec./m ²	1862	1810	1300
² average abundance, thousand spec./m ²	887	961	1294
¹ proportion of injured or dead specimens, %	1.8	2.6	1.9
² proportion of injured or dead specimens, %	0.5	1.5	0.8
¹ Shannon-Weaver index, bit/spec.	1.51	0.66	0.80
² Shannon-Weaver index, bit/spec.	0.96	1.09	1.02
¹ Shannon-Weaver index, bit/mg	1.17	0.60	0.88
² Shannon-Weaver index, bit/mg	0.74	0.73	0.62
¹ average individual masse, mg	7.75	12.8	8.90
² average individual masse, mg	15.5	17.2	13.7
Note: numerals indicate the corresponding areas			

In both areas, the average mass of zooplankton decreased significantly in the period of impact and returned again to the background one or close to background values after completion.

The diversity of zoobenthos varied in the areas ranging from 16 to 26 species. The background was typical of the Caspian Sea species: worms *Hediste diversicolor*, *Oligochaeta gen. sp.*, molluscs *Abra ovata*, *Cerastoderma lamarcki*; crustaceans *Stenocuma gracilis*, *S. graciloides*, *Pontogammarus (Obesogammarus) obesus*, and insect larvae.

The diversity and abundance of zoobenthos at various stages of surveys changed irregularly (Table 3). It can be explained by aggregation of distribution of benthic organisms. Proportion in the total number of benthic cenosis of dead and/or injured specimens was higher during the experimental work, with close values of the index recorded in background conditions and after impact of seismic exploration.

The proportion of dead and/or injured specimens in the populations of benthic invertebrates was highest in close proximity to the pneumatic source (Figure 5). The value of index decreased with the increasing distance from the source of impact, and at the same time, at a distance of 10 m, it remained above background levels.




Figure 5 – Change in proportion of dead and/or injured specimens in the populations of benthic invertebrates at different distances from the pneumatic source

The value of average weight of individual specimens in cenosis was the highest during the experimental tests.

The dynamics of Shannon-Weaver index values across the sites varied irregularly, but in all cases, a deviation of indicator from the background value and from the values obtained during the period after the end of impact was recorded (Figure 6).






Figure 6 – The dynamics of the Shannon-Weaver index values at different stages of the surveys of the impact of seismic acoustic zoobenthos

Fish fauna was presented by 11-13 bento-pelagic species. The composition of background species include dried fish (*Rutilus rutilus*), sprat (*Clupeonella cultriventris*), silverside (*Atherina boyeri caspia*), steers – *Neogobius melanostomus*, *Mesogobius gymnotrachelus macrophth*, *Neogobius caspius*. At the same time, species composition of fish fauna had a high degree of similarity throughout all stages of surveys (Figures 7, 8).



A – before impact, B – during impact, C – after impact

Figure 7 – Dendrogram of similarity in species composition of the fish fauna in area No.1



A – before impact, B – during impact, C – after impact

Figure 8 – Dendrogram of similarity in species composition of the fish fauna in area No.2

The number and variety indicators of bento-pelagic ichthyocenosis during the seismic operations at both areas were lower compared to background levels and stages after impact (Table 4). Changes of ichthyocenosis structure during the seismic survey were accompanied by a relative increase of average weight of individual specimen.

Table 4 – The structural indicators of fish fauna (bento-pelagic communities) of the Northern Caspian Sea at various surveys stages

Indicator	Background conditions	Experimental tests	After impact
¹ number of species	13	8	10
² number of species	8	6	9
¹ average abundance, spec./effort	355	70	312

Indicator	Background conditions	Experimental tests	After impact
² average abundance, spec./effort	213	112	121
¹ Shannon-Weaver index, bit/spec.	1.8	1.2	1.6
² Shannon-Weaver index, bit/spec.	1.45	1.12	1.51
¹ Shannon-Weaver index, bit/mg	1.45	1.13	1.29
² Shannon-Weaver index, bit/mg	1.40	0.98	1.41
¹ average individual masse, mg	6.1	6.9	5.6
² average individual masse, mg	6.3	7.3	6.9
Note: numerals indicate corresponding areas			

The Shannon-Weaver index recorded during seismic surveys was minimal relative to other stages of the surveys (Figure 9).




Figure 9 – Dynamics of Shannon-Weaver index values at different stages of surveys on the effects of seismic operations on the fish fauna

At the stage of experimental tests, in the populations of bento-pelagic fish species the proportion of traumatized specimens decreased with distance from the pneumatic source (Table 5, Figure 10). On the area No.1, the value of this indicator for all experimental stations was higher by more than 1.5 times, compared to the area No.2.

Thus, our experimental and field studies have shown that, for assessing the impact of seismic operations, the relative indicators – the proportion of injured and/or dead specimens, the values of diversity index of Shannon-Weaver and the size of the average individual weight of specimens in hydrocenoses have the greatest significance to the indicator. The maximum proportion of injured and/or dead specimens in the populations of plankton and benthic invertebrates, as well as the deviation of the values of diversity index of Shannon-Weaver and average individual weight in both communities from background values were recorded during the experimental tests in close proximity to the pneumatic source.

Table 5 – Average fish fauna injury during experimental tests

Indicator	Distance from pneumatic source		
	1 meter	5 meters	10 meters
¹ proportion of injured or dead specimens, %	33.3	13.3	0.0
² proportion of injured or dead specimens, %	68.4	39.2	10.5
Note: numerals indicate the corresponding areas			




Figure 10 – Change in the proportion of dead specimens in zoobenthos populations at different distance from the pneumatic source

In the absence of differences in the species composition of fish fauna in various stages of surveys, the minimum diversity and abundance of fish have been observed during the seismic survey. The increase in the average weight of specimens in ichthyocenosis, along with a reduction in numbers during the seismic survey may indicate avoidance of unfavorable zone by younger fishes. This group of aquatic organisms is the most mobile and during the entire seismic surveys can quickly leave the water area with irritating physical influences (noise, vibration, etc.), which is confirmed by the dynamics of structural indicators of ichthyocenoses. Also important the fact that larger specimens that are less susceptible to the effects of seismic acoustic stay in the area of the seismic operations. Similar examples in the behavioral response of fish observed in materials of other experimental studies on assessment of the impact of seismic sources on aquatic organisms [7-9].

Our results enable us to conclude that in conducting seismic surveys, the representatives of benthopelagic ichthyocenosis are subject to an insignificant impact, which is not reflected in terms of their natural life. Planktonic and benthic invertebrates with relatively low levels of mobility are at a significant risk of injury directly in the area affected by pneumatic source.

REFERENCES

- [1] Guidelines on the methods of hydrobiological analysis of surface water and bottom sediments, L.: Gidrometeoizdat, **1983**. 240 p.
- [2] I. Pravdin. Guidelines for fish study, M.: Food industry, **1966**. 377 p.
- [3] Instructions for gathering and primary processing of the Caspian Sea basin aquatic resources and their habitat, Astrakhan: FGUP, CaspNIRKH, **2011**. 233 p.
- [4] Atlas of Invertebrates of the Caspian Sea, M.: Food industry, **1968**. 417 p.
- [5] E. Kazanchev. Fishes of the Caspian Sea (guide), M.: Light and Food industry, **1981**. 168 p.
- [6] N. Kachinsky. Mechanical and Micro-Aggregative Composition of Soil, Methods of Study, M.: USSR Academy of Science, **1958**. 192 p.
- [7] E. Vekilov, E. Kriksunov, Y. Polonsky. Impact of elastic waves from seismic sources on aquatic organisms for marine geophysical exploration // Information and reference text-book, M.: Moscow State University named after M. Lomonosov, **1995**. 64 p.
- [8] A. Vedenev. Analysis of offshore coastal seismic exploration and well drilling impact on migration of salmon at the Sakhalin island, M.: WWF, **2009**. 20 p.
- [9] P. Makarevich. Impact of offshore oil and gas facilities development on pelagic phytocenoses of the Barents Sea. // Herald of MSTU, Vol.16, No.3, **2013**. p. 482.

В.В. Садомский¹, Е.Г. Крупа², И.М. Аминова³

¹Алматы, "SED" ЖШС, Алматы қ., Қазақстан;

²ҚР БҒМ "Зоология институты" РММ, Алматы қ., Қазақстан;

³"Ақжайық" мемлекеттік табиғи резерваты, Атырау қ., Қазақстан

СОЛТҮСТІК КАСПИЙ АКВАТОРИЯСЫНДАҒЫ ГИДРОБИОНТТАРДЫҢ СЕЙСМОАКУСТИКАЛЫҚ ӘСЕРІН ЭКСПЕРИМЕНТТІК ЗЕРТТЕУЛЕР

Аннотация. Мақалада 2012-2013 жылдардың жазғы кезеңінде Солтүстік Каспийдің зоопланктонына, макрозообентосына және ихтиофаунасына сейсмоакустикалық әсер етудің эксперименталдық және табиғи зерттеулерінің материалдары ұсынылған. Сейсmobарлау жұмыстарының әсер етуін бағалау үшін жарақат алған немесе өлген дарақтар үлесінің, Шеннон-Уивердің әртүрлілік индексі мәнінің және дарақтың гидроценоздардағы орташа жеке массасының шамасының ең көп индикаторлық маңыздылығы бар екендігі көрсетілген.

Жарақат алған немесе өлген дарақтардың планктонды және сутүбі омыртқасыздар популяцияларындағы максималды үлесі пневмокөзден 1 м. және 5 м. арақашықтықта тәжірибелі сынақтарды жүргізу кезеңінде тіркелген болатын. Бұл кезеңде екі топтануда фондық шамалардан Шеннон-Уивердің әртүрлілік индексі шамаларының және дарақтың орташа жеке массасының ауытқуы байқалды.

Ихтиофаунаың түрлік құрамында зерттеулерді жүргізудің барлық кезеңдерінде жоғары дәрежесі бар болды. Әртүрліліктің минималды көрсеткіштері (түрлердің саны, Шеннон-Уивер индексінің мәні) және балықтардың саны сейсmobарлау жұмыстарын жүргізу кезінде байқалған болатын. Дарақтардың ихтиоценоздардағы орташа жеке массасының көбеюі, сейсmobарлау жұмыстарын жүргізу кезінде санның азаюымен қатар, кіші жастағы балықтардың қолайсыз аймақтардан аулақ болуын куәландыруы мүмкін.

Тірек сөздер: зоопланктон, зообентос, балықтар, пневматикалық соққы көздері, сейсмикалық барлау жұмыстары.

УДК 502.53/57.087

В.В. Садомский¹, Е.Г. Крупа², И.М. Аминова³

¹ТОО «SED», Алматы, Казахстан;

²РГП на ПХВ «Институт зоологии» КН МОН РК, Алматы, Казахстан;

³Государственный природный резерват «Ақжайық», Атырау, Казахстан

ЭКСПЕРИМЕНТАЛЬНЫЕ ИССЛЕДОВАНИЯ СЕЙСМОАКУСТИЧЕСКОГО ВОЗДЕЙСТВИЯ НА ГИДРОБИОНТОВ СЕВЕРНОГО КАСПИЯ

Аннотация. В статье представлены материалы экспериментальных и натурных исследований сейсмоакустического воздействия на зоопланктон, зообентос и ихтиофауну Северного Каспия в летний период 2012–2013 гг. Показано, что для оценки влияния сейсморазведочных работ наибольшую индикаторную значимость имеют доля травмированных или мертвых особей, значения индекса разнообразия Шеннона-Уивера и величины средней индивидуальной массы особи в гидроценозах. Максимальная доля травмированных и/или мертвых особей в популяциях планктонных и донных беспозвоночных была зафиксирована в период проведения опытных испытаний на расстоянии 1 м и 5 м от пневмоисточника. В этот же период наблюдалось отклонение значений индекса разнообразия Шеннона-Уивера и средней индивидуальной массы особи в обоих сообществах от фоновых значений. Видовой состав ихтиофауны имел высокую степень сходства на всех этапах проведения исследований. Минимальные показатели разнообразия (число видов, значения индекса Шеннона-Уивера) и численности рыб были отмечены во время проведения сейсморазведочных работ. Увеличение средней индивидуальной массы особи в ихтиоценозах, наряду со снижением численности во время проведения сейсморазведочных работ, может свидетельствовать об избегании рыбами младших возрастов неблагоприятной зоны.

Ключевые слова: зоопланктон, зообентос, ихтиофауна, пневмоисточник, сейсморазведочные работы.

CONTENT

Astrophysics

Burtebayev N., Zazulin D.M., Kerimkulov Zh.K., Baktybayev M., Burtebayeva J., Alimov D.K., Nassurlla M. New measurements of differential cross section for elastic scattering process of $^{16}\text{O}(p,p)^{16}\text{O}$ at astrophysical energies.....5

Technical sciences

Poleshchuk O.Kh., Yarkova A.G., Adyrbekova G.M., Zhurhabayeva L. A., Saidakhmetov P.A. Study of the mechanism of the reaction of triazole's formation of using the density functional theory.....11

Kartbayev T.S. Using the neural network technology in solving the tasks of personal identification19

Biology

Ossikbayeva S.O., Orynbayeva Z.S., Tuleukhanov S.T. The mechanism of polyphenolic compounds on prostate cancer.....23

Medicine

Ozhikhenova A.K., Kurakbayev K.K., Karataev M., Ozhikhenov K.A. Monitoring and analysis of bedspace use in day hospitals.....31

Social sciences

Abdrassilov T.K., Kaldybay K.K. Philosophical and ethical values of buddhism.....35

Technical sciences

Krylova Yu.S., Polyakova V.O., Gzgzyan A.M., Lokshin V.N., Kvetnoy I.M. Endometriosis genitalis externa, immunohistochemical diagnosis of endometrium implantation receptivity..... 42

Drozдов A.M., Zhokhov A.L., Yunusov A.A., Yunusova A.A. Solution of the cosmological problem in the approximations. (Part-2)..... 50

Sagatov I.Ye., Kyashnin A.V., Imammyrzaev U.Ye., Danyarov N.B., Nurlan D.T. The direct results of the radical correction of various forms of the atrioventricular septal defect..... 59

Genbach A.A., Jamankulova N.O. Study of heat and mass transfer in capillary-porous cooling systems of a new class of energy thermal installations.....63

Sadomskiy V., Krupa E., Aminova I. Experimental research of seismic-acoustic impact on the north Caspian aquatic organisms 69

Mahmetova N.M., Solonenko V.G., Bekzhanova S.T. The calculation of free oscillations of an anisotropic three-dimensional array of underground structures..... 78

**Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the work described has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the originality detection service Cross Check <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

www.nauka-nanrk.kz

ISSN 2518-1483 (Online), ISSN 2224-5227 (Print)

<http://www.reports-science.kz/index.php/ru/>

Редакторы *М. С. Ахметова, Д. С. Аленов, Т.А. Апендиев, А.Е. Бейсебаева*
Верстка на компьютере *А.М. Кульгинбаевой*

Подписано в печать 10.02.2017.

Формат 60x881/8. Бумага офсетная. Печать – ризограф.

13 п.л. Тираж 2000. Заказ 1.