

ISSN 2518-1483 (Online),
ISSN 2224-5227 (Print)

2017 • 4

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

БАЯНДАМАЛАРЫ

ДОКЛАДЫ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН

REPORTS

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN

ЖУРНАЛ 1944 ЖЫЛДАН ШЫҒА БАСТАҒАН

ЖУРНАЛ ИЗДАЕТСЯ С 1944 г.

PUBLISHED SINCE 1944



Бас редакторы
х.ғ.д., проф., ҚР ҰҒА академигі **М.Ж. Жұрынов**

Редакция алқасы:

Адекенов С.М. проф., академик (Қазақстан) (бас ред. орынбасары)
Боос Э.Г. проф., академик (Қазақстан)
Величкин В.И. проф., корр.-мүшесі (Ресей)
Вольдемар Вуйцик проф. (Польша)
Гончарук В.В. проф., академик (Украина)
Гордиенко А.И. проф., академик (Белорус)
Дука Г. проф., академик (Молдова)
Илолов М.И. проф., академик (Тәжікстан),
Леска Богуслава проф. (Польша),
Локшин В.Н. проф. чл.-корр. (Қазақстан)
Нараев В.Н. проф. (Ресей)
Неклюдов И.М. проф., академик (Украина)
Нур Изура Удзир проф. (Малайзия)
Перни Стефано проф. (Ұлыбритания)
Потапов В.А. проф. (Украина)
Прокопович Полина проф. (Ұлыбритания)
Омбаев А.М. проф. (Қазақстан)
Өтелбаев М.О. проф., академик (Қазақстан)
Садыбеков М.А. проф., корр.-мүшесі (Қазақстан)
Сатаев М.И. проф., корр.-мүшесі (Қазақстан)
Северский И.В. проф., академик (Қазақстан)
Сикорски Марек проф., (Польша)
Рамазанов Т.С. проф., академик (Қазақстан)
Такибаев Н.Ж. проф., академик (Қазақстан), бас ред. орынбасары
Харин С.Н. проф., академик (Қазақстан)
Чечин Л.М. проф., корр.-мүшесі (Қазақстан)
Харун Парлар проф. (Германия)
Энджун Гао проф. (Қытай)
Эркебаев А.Э. проф., академик (Қырғыстан)

«Қазақстан Республикасы Ұлттық ғылым академиясының баяндамалары»
ISSN 2518-1483 (Online),
ISSN 2224-5227 (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» Республикалық қоғамдық бірлестігі (Алматы қ.)
Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде 01.06.2006 ж.
берілген №5540-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылына 6 рет.
Тиражы: 2000 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
http://nauka-nanrk.kz_reports-science.kz

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2017

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Главный редактор
д.х.н., проф., академик НАН РК **М. Ж. Журинов**

Редакционная коллегия:

Адекенов С.М. проф., академик (Казахстан) (зам. гл. ред.)
Боос Э.Г. проф., академик (Казахстан)
Величкин В.И. проф., чл.-корр. (Россия)
Вольдемар Вуйцик проф. (Польша)
Гончарук В.В. проф., академик (Украина)
Гордиенко А.И. проф., академик (Беларусь)
Дука Г. проф., академик (Молдова)
Илолов М.И. проф., академик (Таджикистан),
Леска Богуслава проф. (Польша),
Локшин В.Н. проф. чл.-корр. (Казахстан)
Нараев В.Н. проф. (Россия)
Неклюдов И.М. проф., академик (Украина)
Нур Изура Удзир проф. (Малайзия)
Перни Стефано проф. (Великобритания)
Потапов В.А. проф. (Украина)
Прокопович Полина проф. (Великобритания)
Омбаев А.М. проф. (Казахстан)
Отелбаев М.О. проф., академик (Казахстан)
Садьбеков М.А. проф., чл.-корр. (Казахстан)
Сатаев М.И. проф., чл.-корр. (Казахстан)
Северский И.В. проф., академик (Казахстан)
Сикорски Марек проф., (Польша)
Рамазанов Т.С. проф., академик (Казахстан)
Такибаев Н.Ж. проф., академик (Казахстан), зам. гл. ред.
Харин С.Н. проф., академик (Казахстан)
Чечин Л.М. проф., чл.-корр. (Казахстан)
Харун Парлар проф. (Германия)
Энджун Гао проф. (Китай)
Эркебаев А.Э. проф., академик (Кыргызстан)

«Доклады Национальной академии наук Республики Казахстан»

ISSN 2518-1483 (Online),

ISSN 2224-5227 (Print)

Собственник: Республиканское общественное объединение «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №5540-Ж, выданное 01.06.2006 г.

Периодичность: 6 раз в год.

Тираж: 2000 экземпляров

Адрес редакции: 050010, г.Алматы, ул.Шевченко, 28, ком.218-220, тел. 272-13-19, 272-13-18

<http://nauka-nanrk.kz> reports-science.kz

©Национальная академия наук Республики Казахстан, 2017 г.

Адрес типографии: ИП «Аруна», г.Алматы, ул.Муратбаева, 75

E d i t o r i n c h i e fdoctor of chemistry, professor, academician of NAS RK **M.Zh. Zhurinov****E d i t o r i a l b o a r d :****Adekenov S.M.** prof., academician (Kazakhstan) (deputy editor in chief)**Boos E.G.** prof., academician (Kazakhstan)**Velichkin V.I.** prof., corr. member (Russia)**Voitsik Valdemar** prof. (Poland)**Goncharuk V.V.** prof., academician (Ukraine)**Gordiyenko A.I.** prof., academician (Belarus)**Duka G.** prof., academician (Moldova)**Ilolov M.I.** prof., academician (Tadjikistan),**Leska Boguslava** prof. (Poland),**Lokshin V.N.** prof., corr. member. (Kazakhstan)**Narayev V.N.** prof. (Russia)**Nekludov I.M.** prof., academician (Ukraine)**Nur Izura Udzir** prof. (Malaysia)**Perni Stephano** prof. (Great Britain)**Potapov V.A.** prof. (Ukraine)**Prokopovich Polina** prof. (Great Britain)**Ombayev A.M.** prof. (Kazakhstan)**Otelbayv M.O.** prof., academician (Kazakhstan)**Sadybekov M.A.** prof., corr. member. (Kazakhstan)**Satayev M.I.** prof., corr. member. (Kazakhstan)**Severskyi I.V.** prof., academician (Kazakhstan)**Sikorski Marek** prof., (Poland)**Ramazanov T.S.** prof., academician (Kazakhstan)**Takibayev N.Zh.** prof., academician (Kazakhstan), deputy editor in chief**Kharin S.N.** prof., academician (Kazakhstan)**Chechin L.M.** prof., corr. member. (Kazakhstan)**Kharun Parlar** prof. (Germany)**Endzhun Gao** prof. (China)**Erkebayev A.Ye.** prof., academician (Kyrgyzstan)**Reports of the National Academy of Sciences of the Republic of Kazakhstan.****ISSN 2224-5227****ISSN 2518-1483 (Online),****ISSN 2224-5227 (Print)**

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 5540-Ж, issued 01.06.2006

Periodicity: 6 times a year

Circulation: 2000 copies

Editorial address: 28, Shevchenko str., of.219-220, Almaty, 050010, tel. 272-13-19, 272-13-18,

<http://nauka-nanrk.kz> / reports-science.kz

© National Academy of Sciences of the Republic of Kazakhstan, 2017

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

A.T. Assanova¹, H.A. Ashirbaev², A.P. Sabalakhova³

¹ Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan;

^{2,3} South-Kazakhstan State University after M.O.Auezov, Shymkent, Kazakhstan;

e-mail: anarasanova@list.ru; ashirbaev_1954@mail.ru; sabalakhova@mail.ru

ON THE NONLOCAL PROBLEM FOR A SYSTEM OF THE PARTIAL INTEGRO-DIFFERENTIAL EQUATIONS OF HYPERBOLIC TYPE

Abstract. The nonlocal problem with data on the characteristics for the system of integro-differential equations of hyperbolic type second order is considered. The questions of the existence and uniqueness of a classical solution to the nonlocal problem are studied. The considered problem is reduced to an equivalent nonlocal problem with integral condition by introducing a new unknown function instead of a integral term in the system of equations. The problem with parameter consists of a nonlocal problem for a system of hyperbolic equations with parameter and the integral relation. Algorithms for finding an approximate solution of the equivalent problem with parameter are constructed and the conditions for their convergence are proved. Sufficient conditions for the existence of unique solution to the problem with parameter are established. Conditions of existence of unique classical solution to the nonlocal problem for the system of integro-differential equations of hyperbolic type are obtained in the terms of initial data. Earlier, the method of reduced to an equivalent family of problems for partial differential equations is applied to study of this problem. Sufficient conditions for the existence of unique classical solution of this problem are found in the terms of some matrix compiled by the initial data.

This results are partially supported by grant of the Ministry of education and science of the Republic Kazakhstan, No 0822 / ГФ4.

Key words: nonlocal problem, system of partial integro-differential equations, parameter, algorithm, approximate solution, unique solvability.

1. Introduction. Note that the interest of nonlocal problems for partial integro-differential equations of hyperbolic type has grown. Nonlocal problems are called boundary value problems, in which instead of the classical boundary conditions for the partial integro-differential equations it is given specified combination of values of the unknown function on the boundary of the domain and within it. Boundary conditions are set on the characteristics of the system of hyperbolic equations. The existence and uniqueness of the classical solutions to nonlocal problems for system of hyperbolic integro-differential equations are set. In the present work we consider the system of hyperbolic integro-differential equations of second order in a rectangular domain. Boundary conditions are specified as a combination of values from the required solution and their partial derivatives on first order. We investigate the questions of existence and uniqueness of the classical solution to nonlocal problem for system of hyperbolic integro-differential equations and its applications. For solve to considered problem we use a method of introduction additional functional parameters. The original problem is reduced to an equivalent problem consisting from Goursat problem for system of hyperbolic equations with functional parameters and integral relations. Sufficient conditions of the unique solvability to investigated problem are established in the terms of initial data. Algorithms of finding solution to the nonlocal problem are constructed. The applicability of the obtained results in an optimal control problems are showed.

2. Statement of problem. On the domain $\Omega = [0, T] \times [0, \omega]$ we consider the nonlocal problem with data on the characteristics for system of partial integro-differential equations of hyperbolic type of second order

$$\frac{\partial^2 u}{\partial t \partial x} = A(t, x) \frac{\partial u}{\partial x} + B(t, x) \frac{\partial u}{\partial t} + C(t, x)u + \int_0^\theta \left[K_1(t, \xi) \frac{\partial u(t, \xi)}{\partial \xi} + K_2(t, \xi) \frac{\partial u(t, \xi)}{\partial t} + K_3(t, \xi)u(t, \xi) \right] d\xi + f(t, x), \quad (t, x) \in \Omega, \quad (1)$$

$$u(t, 0) = \psi(t), \quad t \in [0, T], \quad (2)$$

$$P_2(x) \frac{\partial u(0, x)}{\partial x} + P_1(x) \frac{\partial u(t, x)}{\partial t} \Big|_{t=0} + P_0(x)u(0, x) + L_2(x) \frac{\partial u(\theta, x)}{\partial x} + L_1(x) \frac{\partial u(t, x)}{\partial t} \Big|_{t=\theta} + L_0(x)u(\theta, x) + S_2(x) \frac{\partial u(T, x)}{\partial x} + S_1(x) \frac{\partial u(t, x)}{\partial t} \Big|_{t=T} + S_0(x)u(T, x) = \varphi(x), \quad x \in [0, \omega], \quad (3)$$

where $u(t, x) = \text{col}(u_1(t, x), u_2(t, x), \dots, u_n(t, x))$, the $n \times n$ -matrices $A(t, x)$, $B(t, x)$, $C(t, x)$, $K_1(t, x)$, $K_2(t, x)$, $K_3(t, x)$ and n -vector-function $f(t, x)$ are continuous on Ω , the n -vector-function $\psi(t)$ is continuously differentiable on $[0, T]$, the $n \times n$ -matrices $P_i(x)$, $L_i(x)$, $S_i(x)$, and n -vector-function $\varphi(x)$ are continuous on $[0, \omega]$, $i = 0, 1, 2$, $0 \leq \theta \leq \min(T, \omega)$. The initial data satisfy the condition of approval.

A function $u(t, x) \in C(\Omega, R^n)$ having partial derivatives $\frac{\partial u(t, x)}{\partial x} \in C(\Omega, R^n)$, $\frac{\partial u(t, x)}{\partial t} \in C(\Omega, R^n)$, $\frac{\partial^2 u(t, x)}{\partial t \partial x} \in C(\Omega, R^n)$ is called a classical solution to problem (1)-(3) if for all $(t, x) \in \Omega$ it satisfies the system (1), boundary conditions (2) and (3).

In the present paper we investigate the questions of existence and uniqueness of the classical solutions to the nonlocal problem for system of hyperbolic integro-differential equations (1)-(3) and the approaches of constructing its approximate solutions. For this goals, we applied the method of introduction additional functional parameters proposed in [1-18] for the solve of nonlocal boundary value problems for systems of hyperbolic equations with mixed derivative. Considered problem is provided to nonlocal problem with integral condition for system of hyperbolic equations including additional function. Hence, this problem is reduced to an equivalent problem, consisting of Goursat problem for the system of hyperbolic equations with functional parameters and Cauchy problem for system of ordinary differential equations with respect to the entered parameters by introducing new unknown functions. The algorithm of finding the approximate solution of the investigated problem is proposed and its convergence proved. Sufficient conditions of the existence of unique classical solution to problem (1)-(3) are obtained in the terms of initial data. The applicability of the obtained results in the optimal control problems is showed.

3. *Reduction to nonlocal problem with integral condition for system of hyperbolic equations including special function.* We introduce an additional special function

$$\mu(t) = \int_0^\theta \left[K_1(t, \xi) \frac{\partial u(t, \xi)}{\partial \xi} + K_2(t, \xi) \frac{\partial u(t, \xi)}{\partial t} + K_3(t, \xi)u(t, \xi) \right] d\xi$$

and write the problem (1)-(3) in the following form

$$\frac{\partial^2 u}{\partial t \partial x} = A(t, x) \frac{\partial u}{\partial x} + B(t, x) \frac{\partial u}{\partial t} + C(t, x)u + \mu(t) + f(t, x), \quad (t, x) \in \Omega, \quad (4)$$

$$u(t, 0) = \psi(t), \quad t \in [0, T], \quad (5)$$

$$P_2(x) \frac{\partial u(0, x)}{\partial x} + P_1(x) \frac{\partial u(t, x)}{\partial t} \Big|_{t=0} + P_0(x) u(0, x) + L_2(x) \frac{\partial u(\theta, x)}{\partial x} + L_1(x) \frac{\partial u(t, x)}{\partial t} \Big|_{t=\theta} + \\ + L_0(x) u(\theta, x) + S_2(x) \frac{\partial u(T, x)}{\partial x} + S_1(x) \frac{\partial u(t, x)}{\partial t} \Big|_{t=T} + S_0(x) u(T, x) = \varphi(x), \quad x \in [0, \omega], \quad (6)$$

$$\mu(t) = \int_0^\theta \left[K_1(t, \xi) \frac{\partial u(t, \xi)}{\partial \xi} + K_2(t, \xi) \frac{\partial u(t, \xi)}{\partial t} + K_3(t, \xi) u(t, \xi) \right] d\xi, \quad t \in [0, T]. \quad (7)$$

4. *Scheme of the method and algorithm.* Let $\lambda(x) = u(0, x)$. In the problem (4)-(7) we change the function $u(t, x)$ by $u(t, x) = \tilde{u}(t, x) + \lambda(x)$ and proceed to the following equivalent problem

$$\frac{\partial^2 \tilde{u}}{\partial t \partial x} = A(t, x) \frac{\partial \tilde{u}}{\partial x} + B(t, x) \frac{\partial \tilde{u}}{\partial t} + C(t, x) \tilde{u} + A(t, x) \dot{\lambda}(x) + C(t, x) \lambda(x) + \mu(t) + f(t, x), \quad (8)$$

$$\tilde{u}(t, 0) + \lambda(0) = \psi(t), \quad t \in [0, T], \quad (9)$$

$$\tilde{u}(0, x) = 0, \quad x \in [0, \omega], \quad (10)$$

$$[P_2(x) + L_2(x) + S_2(x)] \dot{\lambda}(x) + [P_0(x) + L_0(x) + S_0(x)] \lambda(x) + \\ + P_1(x) \frac{\partial \tilde{u}(t, x)}{\partial t} \Big|_{t=0} + L_2(x) \frac{\partial \tilde{u}(\theta, x)}{\partial x} + L_1(x) \frac{\partial \tilde{u}(t, x)}{\partial t} \Big|_{t=\theta} + L_0(x) \tilde{u}(\theta, x) + S_2(x) \frac{\partial \tilde{u}(T, x)}{\partial x} + \\ + S_1(x) \frac{\partial \tilde{u}(t, x)}{\partial t} \Big|_{t=T} + S_0(x) \tilde{u}(T, x) = \varphi(x), \quad x \in [0, \omega]. \quad (11)$$

$$\mu(t) = \int_0^\theta \left[K_1(t, \xi) \frac{\partial \tilde{u}(t, \xi)}{\partial \xi} + K_2(t, \xi) \frac{\partial \tilde{u}(t, \xi)}{\partial t} + K_3(t, \xi) \tilde{u}(t, \xi) + K_1(t, \xi) \dot{\lambda}(\xi) + K_3(t, \xi) \lambda(\xi) \right] d\xi. \quad (12)$$

A triple functions $(\tilde{u}(t, x), \lambda(x), \mu(t))$, where the function $\tilde{u}(t, x) \in C(\Omega, R^n)$ has partial derivatives $\frac{\partial \tilde{u}(t, x)}{\partial x} \in C(\Omega, R^n)$, $\frac{\partial \tilde{u}(t, x)}{\partial t} \in C(\Omega, R^n)$, $\frac{\partial^2 \tilde{u}(t, x)}{\partial t \partial x} \in C(\Omega, R^n)$, the function $\lambda(x) \in C([0, \omega], R^n)$ has derivative $\dot{\lambda}(x) \in C([0, \omega], R^n)$, the function $\mu(t) \in C([0, T], R^n)$ determine from relations (12) for all $t \in [0, T]$, is called a solution to problem (8)-(12) if for all $(t, x) \in \Omega$ it satisfies of the system of hyperbolic equations with parameters (8), the boundary conditions (9), (10), the functional relation (11) and the integral condition (12).

From the compatibility condition at the point (0,0) of initial data is yield: $\lambda(0) = \psi(0)$. Then the condition (9) may be rewrite in the following form

$$\tilde{u}(t, 0) = \psi(t) - \psi(0), \quad t \in [0, T]. \quad (13)$$

The problem (8), (10), (13) at fixed $\lambda(x)$, $\mu(t)$ is the Goursat problem with respect to $\tilde{u}(t, x)$ on Ω . The relation (11) allows us to determine the unknown functional parameter $\lambda(x)$. The integral condition (12) allows us to determine the unknown function $\mu(t)$ for all $t \in [0, T]$.

We introduce new unknown functions $\tilde{v}(t, x) = \frac{\partial \tilde{u}(t, x)}{\partial x}$, $\tilde{w}(t, x) = \frac{\partial \tilde{u}(t, x)}{\partial t}$. Goursat problem (8), (10), (13) is equivalent to a three systems integral equations

$$\begin{aligned} \tilde{v}(t, x) = & \int_0^t \{A(\tau, x)\tilde{v}(\tau, x) + B(\tau, x)\tilde{w}(\tau, x) + C(\tau, x)\tilde{u}(\tau, x) + \\ & + A(\tau, x)\dot{\lambda}(x) + C(\tau, x)\lambda(x) + \mu(\tau) + f(\tau, x)\}d\tau, \end{aligned} \quad (14)$$

$$\begin{aligned} \tilde{w}(t, x) = & \psi(t) + \int_0^x \{A(t, \xi)\tilde{v}(t, \xi) + B(t, \xi)\tilde{w}(t, \xi) + C(t, \xi)\tilde{u}(t, \xi) + \\ & + A(t, \xi)\dot{\lambda}(\xi) + C(t, \xi)\lambda(\xi) + \mu(t) + f(t, \xi)\}d\xi, \end{aligned} \quad (15)$$

$$\begin{aligned} \tilde{u}(t, x) = & \psi(t) - \psi(0) + \int_0^t \int_0^x \{A(\tau, \xi)\tilde{v}(\tau, \xi) + B(\tau, \xi)\tilde{w}(\tau, \xi) + C(\tau, \xi)\tilde{u}(\tau, \xi) + \\ & + A(\tau, \xi)\dot{\lambda}(\xi) + C(\tau, \xi)\lambda(\xi) + \mu(\tau) + f(\tau, \xi)\}d\xi d\tau. \end{aligned} \quad (16)$$

In the relation (11) instead of the functions $\tilde{v}(\theta, x)$, $\tilde{v}(T, x)$, we substitute the appropriate expressions of the integral relation under $t = \theta$, $t = T$, respectively. Then we obtain

$$\begin{aligned} & \left[P_2(x) + L_2(x) + L_2(x) \int_0^\theta A(\tau, x) d\tau + S_2(x) + S_2(x) \int_0^T A(\tau, x) d\tau \right] \dot{\lambda}(x) = \\ & = - \left[P_0(x) + L_0(x) + S_0(x) + L_2(x) \int_0^\theta C(\tau, x) d\tau + S_2(x) \int_0^T C(\tau, x) d\tau \right] \lambda(x) - \\ & - L_2(x) \int_0^\theta \mu(\tau) d\tau - S_2(x) \int_0^T \mu(\tau) d\tau - P_1(x)\tilde{w}(0, x) - L_1(x)\tilde{w}(\theta, x) - S_1(x)\tilde{w}(T, x) - \\ & - L_2(x) \int_0^\theta \{A(\tau, x)\tilde{v}(\tau, x) + B(\tau, x)\tilde{w}(\tau, x) + C(\tau, x)\tilde{u}(\tau, x)\}d\tau - L_0(x)\tilde{u}(\theta, x) - \\ & - S_2(x) \int_0^T \{A(\tau, x)\tilde{v}(\tau, x) + B(\tau, x)\tilde{w}(\tau, x) + C(\tau, x)\tilde{u}(\tau, x)\}d\tau - S_0(x)\tilde{u}(T, x) - \\ & - L_2(x) \int_0^\theta f(\tau, x) d\tau - S_2(x) \int_0^T f(\tau, x) d\tau + \varphi(x), \quad x \in [0, \omega]. \end{aligned} \quad (17)$$

From the compatibility condition follows the initial condition

$$\lambda(0) = \psi(0). \quad (18)$$

The unknown functional parameter $\lambda(x)$ will be determined from Cauchy problem for system of ordinary differential equations (17), (18). The unknown special function $\mu(t)$ will be determined from integral relation (12).

If we know the functional parameter $\lambda(x)$, the special function $\mu(t)$, then from integral systems (14)-(16) find the functions $\tilde{u}(t, x)$, $\tilde{v}(t, x)$, $\tilde{w}(t, x)$. Conversely, if we know functions $\tilde{u}(t, x)$, $\tilde{v}(t, x)$, $\tilde{w}(t, x)$, then from Cauchy problem (17), (18) and integral condition (12) we find the functional parameter $\lambda(x)$ and special function $\mu(t)$. Since the functions $\tilde{u}(t, x)$, $\tilde{v}(t, x)$, $\tilde{w}(t, x)$ and $\lambda(x)$, $\mu(t)$ are unknown together for finding of the solution to problem (8)--(12) we use an iterative method. The solution to problem (8)--(12) is the triple functions $(\tilde{u}^*(t, x), \lambda^*(x), \mu^*(t))$ we defined as a limit of sequence of triples $(\tilde{u}^{(m)}(t, x), \lambda^{(m)}(x), \mu^{(m)}(t, x))$, $m = 0, 1, 2, \dots$, according to the following algorithm:

Step 0. 1) Let the matrix $D_1(x) = P_2(x) + L_2(x) + L_2(x) \int_0^\theta A(\tau, x) d\tau + S_2(x) + S_2(x) \int_0^T A(\tau, x) d\tau$ is invertible for all $x \in [0, \omega]$. Suppose in the right-hand part of the system (17) $\mu(t) = 0$, $\tilde{u}(t, x) = \psi(t) - \psi(0)$, $\tilde{v}(t, x) = 0$, $\tilde{w}(t, x) = \dot{\psi}(t)$, from Cauchy problem (17), (18) we find the initial approximation $\lambda^{(0)}(x)$ for all $x \in [0, \omega]$:

$$\begin{aligned} \lambda^{(0)}(x) = & \psi(0) - \int_0^x D_1^{-1}(\xi) D_2(\xi) \lambda^{(0)}(\xi) d\xi - \\ & - \int_0^x D_1^{-1}(\xi) [P_1(\xi) \dot{\psi}(0) + L_1(\xi) \dot{\psi}(\theta) + S_1(\xi) \dot{\psi}(T) + L_0(\xi) \{\psi(\theta) - \psi(0)\} + S_0(\xi) \{\psi(T) - \psi(0)\}] d\xi - \\ & - \int_0^x D_1^{-1}(\xi) L_2(\xi) \int_0^\theta \{B(\tau, \xi) \dot{\psi}(\tau) + C(\tau, \xi) [\psi(\tau) - \psi(0)]\} d\tau d\xi - \\ & - \int_0^x D_1^{-1}(\xi) S_2(\xi) \int_0^T \{B(\tau, \xi) \dot{\psi}(\tau) + C(\tau, \xi) [\psi(\tau) - \psi(0)]\} d\tau d\xi - \\ & - \int_0^x D_1^{-1}(\xi) \left[L_2(\xi) \int_0^\theta f(\tau, \xi) d\tau + S_2(\xi) \int_0^T f(\tau, \xi) d\tau - \varphi(\xi) \right] d\xi, \end{aligned}$$

where $D_2(x) = P_0(x) + L_0(x) + S_0(x) + L_2(x) \int_0^\theta C(\tau, x) d\tau + S_2(x) \int_0^T C(\tau, x) d\tau$.

2) From the system of integral equations (14)--(16) under $\mu(t) = 0$, $\lambda(x) = \lambda^{(0)}(x)$, $\dot{\lambda}(x) = \dot{\lambda}^{(0)}(x)$ we find the functions $\tilde{u}^{(0)}(t, x)$, $\tilde{v}^{(0)}(t, x)$, $\tilde{w}^{(0)}(t, x)$ for all $(t, x) \in \Omega$:

$$\begin{aligned} \tilde{v}^{(0)}(t, x) = & \int_0^t \{A(\tau, x) \tilde{v}^{(0)}(\tau, x) + B(\tau, x) \tilde{w}^{(0)}(\tau, x) + C(\tau, x) \tilde{u}^{(0)}(\tau, x) + \\ & + A(\tau, x) \dot{\lambda}^{(0)}(x) + C(\tau, x) \lambda^{(0)}(x) + f(\tau, x)\} d\tau, \\ \tilde{w}^{(0)}(t, x) = & \dot{\psi}(t) + \int_0^x \{A(t, \xi) \tilde{v}^{(0)}(t, \xi) + B(t, \xi) \tilde{w}^{(0)}(t, \xi) + C(t, \xi) \tilde{u}^{(0)}(t, \xi) + \\ & + A(t, \xi) \dot{\lambda}^{(0)}(\xi) + C(t, \xi) \lambda^{(0)}(\xi) + f(t, \xi)\} d\xi, \\ \tilde{u}^{(0)}(t, x) = & \psi(t) - \psi(0) + \int_0^t \int_0^x \{A(\tau, \xi) \tilde{v}^{(0)}(\tau, \xi) + B(\tau, \xi) \tilde{w}^{(0)}(\tau, \xi) + C(\tau, \xi) \tilde{u}^{(0)}(\tau, \xi) + \\ & + A(\tau, \xi) \dot{\lambda}^{(0)}(\xi) + C(\tau, \xi) \lambda^{(0)}(\xi) + f(\tau, \xi)\} d\xi d\tau. \end{aligned}$$

From integral relation (12) under $\tilde{u}(t, x) = \tilde{u}^{(0)}(t, x)$, $\tilde{v}(t, x) = \tilde{v}^{(0)}(t, x)$, $\tilde{w}(t, x) = \tilde{w}^{(0)}(t, x)$, $\lambda(x) = \lambda^{(0)}(x)$, $\dot{\lambda}(x) = \dot{\lambda}^{(0)}(x)$ we find the initial approximation $\mu^{(0)}(t)$ for all $t \in [0, T]$:

$$\mu^{(0)}(t) = \int_0^\theta [K_1(t, \xi) \tilde{v}^{(0)}(t, \xi) + K_2(t, \xi) \tilde{w}^{(0)}(t, \xi) + K_3(t, \xi) \tilde{u}^{(0)}(t, \xi) + K_1(t, \xi) \dot{\lambda}^{(0)}(\xi) + K_3(t, \xi) \lambda^{(0)}(\xi)] d\xi.$$

Step 1. 1) Suppose in the right-hand part of the system (17) $\mu(t) = \mu^{(0)}(t)$, $\tilde{u}(t, x) = \tilde{u}^{(0)}(t, x)$, $\tilde{v}(t, x) = \tilde{v}^{(0)}(t, x)$, $\tilde{w}(t, x) = \tilde{w}^{(0)}(t, x)$, from Cauchy problem (17), (18) we find the first approximation $\lambda^{(1)}(x)$ for all $x \in [0, \omega]$:

$$\begin{aligned} \lambda^{(1)}(x) = & \psi(0) - \int_0^x D_1^{-1}(\xi) D_2(\xi) \lambda^{(1)}(\xi) d\xi - \\ & - \int_0^x D_1^{-1}(\xi) L_2(\xi) d\xi \int_0^\theta \mu^{(0)}(\tau) d\tau - \int_0^x D_1^{-1}(\xi) S_2(\xi) d\xi \int_0^T \mu^{(0)}(\tau) d\tau - \\ & - \int_0^x D_1^{-1}(\xi) [P_1(\xi) \tilde{w}^{(0)}(0, \xi) + L_1(\xi) \tilde{w}^{(0)}(\theta, \xi) + S_1(\xi) \tilde{w}^{(0)}(T, \xi) + L_0(\xi) \tilde{u}^{(0)}(\theta, \xi) + S_0(\xi) \tilde{u}^{(0)}(T, \xi)] d\xi - \\ & - \int_0^x D_1^{-1}(\xi) L_2(\xi) \int_0^\theta \{A(\tau, \xi) \tilde{v}^{(0)}(\tau, \xi) B(\tau, \xi) \tilde{w}^{(0)}(\tau, \xi) + C(\tau, \xi) \tilde{u}^{(0)}(\tau, \xi)\} d\tau d\xi - \\ & - \int_0^x D_1^{-1}(\xi) S_2(\xi) \int_0^T \{A(\tau, \xi) \tilde{v}^{(0)}(\tau, \xi) B(\tau, \xi) \tilde{w}^{(0)}(\tau, \xi) + C(\tau, \xi) \tilde{u}^{(0)}(\tau, \xi)\} d\tau d\xi - \\ & - \int_0^x D_1^{-1}(\xi) \left[L_2(\xi) \int_0^\theta f(\tau, \xi) d\tau + S_2(\xi) \int_0^T f(\tau, \xi) d\tau - \varphi(\xi) \right] d\xi. \end{aligned}$$

2) From the system of integral equations (14)–(16) under $\mu(t) = \mu^{(0)}(t)$, $\lambda(x) = \lambda^{(1)}(x)$, $\dot{\lambda}(x) = \dot{\lambda}^{(1)}(x)$ we find the functions $\tilde{u}^{(1)}(t, x)$, $\tilde{v}^{(1)}(t, x)$, $\tilde{w}^{(1)}(t, x)$ for all $(t, x) \in \Omega$:

$$\begin{aligned} \tilde{v}^{(1)}(t, x) = & \int_0^t \{A(\tau, x) \tilde{v}^{(1)}(\tau, x) + B(\tau, x) \tilde{w}^{(1)}(\tau, x) + C(\tau, x) \tilde{u}^{(1)}(\tau, x) + \\ & + A(\tau, x) \dot{\lambda}^{(1)}(x) + C(\tau, x) \lambda^{(1)}(x) + \mu^{(0)}(\tau) + f(\tau, x)\} d\tau, \\ \tilde{w}^{(1)}(t, x) = & \psi(t) + \int_0^x \{A(t, \xi) \tilde{v}^{(1)}(t, \xi) + B(t, \xi) \tilde{w}^{(1)}(t, \xi) + C(t, \xi) \tilde{u}^{(1)}(t, \xi) + \\ & + A(t, \xi) \dot{\lambda}^{(1)}(\xi) + C(t, \xi) \lambda^{(1)}(\xi) + \mu^{(0)}(t) + f(t, \xi)\} d\xi, \\ \tilde{u}^{(1)}(t, x) = & \psi(t) - \psi(0) + \int_0^t \int_0^x \{A(\tau, \xi) \tilde{v}^{(1)}(\tau, \xi) + B(\tau, \xi) \tilde{w}^{(1)}(\tau, \xi) + C(\tau, \xi) \tilde{u}^{(1)}(\tau, \xi) + \\ & + A(\tau, \xi) \dot{\lambda}^{(1)}(\xi) + C(\tau, \xi) \lambda^{(1)}(\xi) + \mu^{(0)}(\tau) + f(\tau, \xi)\} d\xi d\tau. \end{aligned}$$

From integral relation (12) under $\tilde{u}(t, x) = \tilde{u}^{(1)}(t, x)$, $\tilde{v}(t, x) = \tilde{v}^{(1)}(t, x)$, $\tilde{w}(t, x) = \tilde{w}^{(1)}(t, x)$, $\lambda(x) = \lambda^{(1)}(x)$, $\dot{\lambda}(x) = \dot{\lambda}^{(1)}(x)$ we find the first approximation $\mu^{(1)}(t)$ for all $t \in [0, T]$:

$$\mu^{(1)}(t) = \int_0^\theta [K_1(t, \xi) \tilde{v}^{(1)}(t, \xi) + K_2(t, \xi) \tilde{w}^{(1)}(t, \xi) + K_3(t, \xi) \tilde{u}^{(1)}(t, \xi) + K_1(t, \xi) \dot{\lambda}^{(1)}(\xi) + K_3(t, \xi) \lambda^{(1)}(\xi)] d\xi.$$

And so on.

Step m . 1) Suppose in the right-hand part of the system (17) $\mu(t) = \mu^{(m-1)}(t)$, $\tilde{u}(t, x) = \tilde{u}^{(m-1)}(t, x)$, $\tilde{v}(t, x) = \tilde{v}^{(m-1)}(t, x)$, $\tilde{w}(t, x) = \tilde{w}^{(m-1)}(t, x)$, from Cauchy problem (17), (18) we find the m -th approximation $\lambda^{(m)}(x)$ for all $x \in [0, \omega]$:

$$\begin{aligned} \lambda^{(m)}(x) = & \psi(0) - \int_0^x D_1^{-1}(\xi) D_2(\xi) \lambda^{(m)}(\xi) d\xi - \int_0^x D_1^{-1}(\xi) \left[L_2(\xi) \int_0^\theta \mu^{(m-1)}(\tau) d\tau + S_2(\xi) \int_0^T \mu^{(m-1)}(\tau) d\tau \right] d\xi - \\ & - \int_0^x D_1^{-1}(\xi) [P_1(\xi) \tilde{w}^{(m-1)}(0, \xi) + L_1(\xi) \tilde{w}^{(m-1)}(\theta, \xi) + S_1(\xi) \tilde{w}^{(m-1)}(T, \xi) + L_0(\xi) \tilde{u}^{(m-1)}(\theta, \xi) + S_0(\xi) \tilde{u}^{(m-1)}(T, \xi)] d\xi - \end{aligned}$$

$$\begin{aligned}
 & - \int_0^x D_1^{-1}(\xi) L_2(\xi) \int_0^\theta \left\{ A(\tau, \xi) \tilde{v}^{(m-1)}(\tau, \xi) + B(\tau, \xi) \tilde{w}^{(m-1)}(\tau, \xi) + C(\tau, \xi) \tilde{u}^{(m-1)}(\tau, \xi) \right\} d\tau d\xi - \\
 & - \int_0^x D_1^{-1}(\xi) S_2(\xi) \int_0^T \left\{ A(\tau, \xi) \tilde{v}^{(m-1)}(\tau, \xi) + B(\tau, \xi) \tilde{w}^{(m-1)}(\tau, \xi) + C(\tau, \xi) \tilde{u}^{(m-1)}(\tau, \xi) \right\} d\tau d\xi - \\
 & - \int_0^x D_1^{-1}(\xi) \left[L_2(\xi) \int_0^\theta f(\tau, \xi) d\tau + S_2(\xi) \int_0^T f(\tau, \xi) d\tau - \varphi(\xi) \right] d\xi, \quad x \in [0, \omega].
 \end{aligned}$$

2) From the system of integral equations (14)--(16) under $\mu(t) = \mu^{(m-1)}(t)$, $\lambda(x) = \lambda^{(m)}(x)$, $\dot{\lambda}(x) = \dot{\lambda}^{(m)}(x)$ we find the functions $\tilde{u}^{(m)}(t, x)$, $\tilde{v}^{(m)}(t, x)$, $\tilde{w}^{(m)}(t, x)$ for all $(t, x) \in \Omega$:

$$\begin{aligned}
 \tilde{v}^{(m)}(t, x) &= \int_0^t \left\{ A(\tau, x) \tilde{v}^{(m)}(\tau, x) + B(\tau, x) \tilde{w}^{(m)}(\tau, x) + C(\tau, x) \tilde{u}^{(m)}(\tau, x) + \right. \\
 & \left. + A(\tau, x) \dot{\lambda}^{(m)}(x) + C(\tau, x) \lambda^{(m)}(x) + \mu^{(m-1)}(\tau) + f(\tau, x) \right\} d\tau, \\
 \tilde{w}^{(m)}(t, x) &= \psi(t) + \int_0^x \left\{ A(t, \xi) \tilde{v}^{(m)}(t, \xi) + B(t, \xi) \tilde{w}^{(m)}(t, \xi) + C(t, \xi) \tilde{u}^{(m)}(t, \xi) + \right. \\
 & \left. + A(t, \xi) \dot{\lambda}^{(m)}(\xi) + C(t, \xi) \lambda^{(m)}(\xi) + \mu^{(m-1)}(t) + f(t, \xi) \right\} d\xi, \\
 \tilde{u}^{(m)}(t, x) &= \psi(t) - \psi(0) + \int_0^t \int_0^x \left\{ A(\tau, \xi) \tilde{v}^{(m)}(\tau, \xi) + B(\tau, \xi) \tilde{w}^{(m)}(\tau, \xi) + C(\tau, \xi) \tilde{u}^{(m)}(\tau, \xi) + \right. \\
 & \left. + A(\tau, \xi) \dot{\lambda}^{(m)}(\xi) + C(\tau, \xi) \lambda^{(m)}(\xi) + \mu^{(m-1)}(\tau) + f(\tau, \xi) \right\} d\xi d\tau.
 \end{aligned}$$

From integral relation (12) under $\tilde{u}(t, x) = \tilde{u}^{(m)}(t, x)$, $\tilde{v}(t, x) = \tilde{v}^{(m)}(t, x)$, $\tilde{w}(t, x) = \tilde{w}^{(m)}(t, x)$, $\lambda(x) = \lambda^{(m)}(x)$, $\dot{\lambda}(x) = \dot{\lambda}^{(m)}(x)$ we find the m -th approximation $\mu^{(m)}(t)$ for all $t \in [0, T]$:

$$\begin{aligned}
 \mu^{(m)}(t) &= \int_0^\theta \left[K_1(t, \xi) \tilde{v}^{(m)}(t, \xi) + K_2(t, \xi) \tilde{w}^{(m)}(t, \xi) + K_3(t, \xi) \tilde{u}^{(m)}(t, \xi) \right] d\xi + \\
 & + \int_0^\theta \left[K_1(t, \xi) \dot{\lambda}^{(m)}(\xi) + K_3(t, \xi) \lambda^{(m)}(\xi) \right] d\xi, \quad m = 1, 2, 3, \dots
 \end{aligned}$$

5. The main result.

Let $a = \max_{(t,x) \in \Omega} \|A(t, x)\|$, $b = \max_{(t,x) \in \Omega} \|B(t, x)\|$, $c = \max_{(t,x) \in \Omega} \|C(t, x)\|$, $H = a + b + c$,

$\alpha_0 = \max_{x \in [0, \omega]} \| [D_1(x)]^{-1} \|$, $\alpha = \max_{x \in [0, \omega]} \| [D_1(x)]^{-1} D_2(x) \|$,

$a_1 = \max_{x \in [0, \omega]} \|P_1(x)\| + \max_{x \in [0, \omega]} \|L_1(x)\| + \max_{x \in [0, \omega]} \|S_1(x)\| + \max_{x \in [0, \omega]} \|L_0(x)\| + \max_{x \in [0, \omega]} \|S_0(x)\|$,

$b_1 = \max_{x \in [0, \omega]} \|L_2(x)\| \max(T, \omega) \left[e^{H(\theta+\omega)} - e^{H\omega} \right] + \max_{x \in [0, \omega]} \|S_2(x)\| \max(T, \omega) \left[e^{H(T+\omega)} - e^{H\omega} \right]$,

$a_2 = \theta \max_{x \in [0, \omega]} \|L_2(x)\| + T \max_{x \in [0, \omega]} \|S_2(x)\|$, $b_2 = \max(T, \omega) e^{H(T+\omega)} (a + c + 1)$,

$c_1 = \theta \left[(b_2 + 1) \left(\max_{(t,x) \in \Omega} \|K_1(t, x)\| + \max_{(t,x) \in \Omega} \|K_3(t, x)\| \right) + b_2 \max_{(t,x) \in \Omega} \|K_2(t, x)\| \right]$.

The following theorem gives conditions of realizability and convergence of the constructed algorithm and the conditions of the existence of unique solution to problem (8)--(12).

Theorem 1. Suppose that

i) the matrix $D_1(x)$ is invertible for all $x \in [0, \omega]$;

ii) the inequality fulfilled

$$q(T, \omega) = \max([\alpha e^{\alpha\omega} \omega + 1]\alpha_0(a_2c_1 + (a_1 + b_1)b_2), e^{\alpha\omega} \omega \alpha_0(a_2c_1 + (a_1 + b_1)b_2), c_1) < 1.$$

Then the problem for system of hyperbolic equations with parameters (8)–(12) has a unique solution.

Theorem 2. Suppose that the conditions i) - ii) of Theorem 1 are fulfilled.

Then the nonlocal problem for system of partial integro-differential equations (1)–(3) has a unique classical solution.

The proof of the theorem is similar to the scheme of the proof of theorem [12, p. 26].

REFERENCES

- [1] Asanova A.T., Dzhumabaev D.S. Unique solvability of the boundary value problem for systems of hyperbolic equations with data on the characteristics // Computational Mathematics and Mathematical Physics. **2002**. Vol. 42. No 11. P. 1609-1621.
- [2] Asanova A.T., Dzhumabaev D.S. Correct solvability of a nonlocal boundary value problem for systems of hyperbolic equations // Doklady Mathematics. **2003**. Vol. 68. No 1. P. 46-49.
- [3] Asanova A.T., Dzhumabaev D.S. Unique solvability of nonlocal boundary value problems for systems of hyperbolic equations // Differential Equations. **2003**. Vol. 39. No 10. P. 1414-1427.
- [4] Asanova A.T., Dzhumabaev D.S. Periodic solutions of systems of hyperbolic equations bounded on a plane // Ukrainian Mathematical Journal. **2004**. Vol. 56. No 4. P. 682-694.
- [5] Asanova A.T., Dzhumabaev D.S. Well-posed solvability of nonlocal boundary value problems for systems of hyperbolic equations // Differential Equations. **2005**. Vol. 41. No 3. P. 352-363.
- [6] Asanova A.T. A nonlocal boundary value problem for systems of quasilinear hyperbolic equations // Doklady Mathematics. **2006**. Vol. 74. No 3. P. 787-791.
- [7] Asanova A.T. On the unique solvability of a family of two-point boundary-value problems for systems of ordinary differential equations // Journal of Mathematical Sciences. **2008**. Vol. 150. No 5. P. 2302-2316.
- [8] Asanova A.T. On the unique solvability of a nonlocal boundary value problem with data on intersecting lines for systems of hyperbolic equations // Differential Equations. **2009**. Vol. 45. No 3. P. 385-394.
- [9] Asanova A.T. On a boundary-value problem with data on noncharacteristic intersecting lines for systems of hyperbolic equations with mixed derivative // Journal of Mathematical Sciences (United States). **2012**. Vol. 187. No 4. P. 375-386.
- [10] Asanova A.T. On a nonlocal boundary-value problem for systems of impulsive hyperbolic equations // Ukrainian Mathematical Journal. **2013**. Vol. 65. No 3. P. 349-365.
- [11] Asanova A.T., Dzhumabaev D.S. Well-posedness of nonlocal boundary value problems with integral condition for the system of hyperbolic equations // Journal of Mathematical Analysis and Applications. **2013**. Vol. 402. No 1. P. 167-178.
- [12] Asanova A.T. On a solvability of the nonlocal problem with integral conditions for system of the equations of hyperbolic type // Mathematical journal. **2014**. Vol. 14. No 2 (52). P. 21-35. [in Russian]
- [13] Asanova A.T. Well-posed solvability of a nonlocal boundary-value problem for systems of hyperbolic equations with impulse effects // Ukrainian Mathematical Journal. **2015**. Vol. 67. No 3. P. 333-346.
- [14] Asanova A.T. On solvability of nonlinear boundary value problems with integral condition for the system of hyperbolic equations // Electronic Journal of Qualitative Theory of Differential Equations. **2015**. No 63. P. 1-13.
- [15] Asanova A.T., Imanchiev A.E. On conditions of the solvability of nonlocal multi-point boundary value problems for quasi-linear systems of hyperbolic equations // Eurasian Mathematical Journal. **2015**. Vol. 6. No 4. P. 19-28.
- [16] Asanova A.T. Multipoint problem for a system of hyperbolic equations with mixed derivative // Journal of Mathematical Sciences (United States). **2016**. Vol. 212. No 3. P. 213-233.
- [17] Asanova A.T. Criteria of solvability of nonlocal boundary-value problem for systems of hyperbolic equations with mixed derivatives // Russian Mathematics. **2016**. Vol. 60. No 1. P. 1-17.
- [18] Asanova A.T. On the solvability of nonlocal boundary value problem for the systems of impulsive hyperbolic equations with mixed derivatives // Journal of Discontinuity, Nonlinearity and Complexity. **2016**. Vol. 5. No 2. P. 153-165.

А.Т. Асанова¹, Х.А. Аширбаев², А.П. Сабалахова³

¹Математика және математикалық модельдеу институты, Алматы қ., Қазақстан;

^{2,3}М. Әуезов атындағы Оңтүстік Қазақстан мемлекеттік университеті, Шымкент қ., Қазақстан

ГИПЕРБОЛАЛЫҚ ТЕКТЕС ДЕРБЕС ТУЫНДЫЛЫ ИНТЕГРАЛДЫҚ-ДИФФЕРЕНЦИАЛДЫҚ ТЕНДЕУЛЕР ЖҮЙЕСІ ҮШІН БЕЙЛОКАЛ ЕСЕП ТУРАЛЫ

Аннотация. Екінші ретті гиперболалық тектес интегралдық-дифференциалдық тендеулер жүйесі үшін бейлокал есеп қарастырылады. Бейлокал есептің классикалық шешімінің бар болуы мен жалғыздығы мәселелері зерттелген. Интегралдық қосылғыштың орнына жаңа белгісіз функция енгізу жолымен зерттеліп отырған есеп пара-пара интегралдық шарты бар бейлокал есепке келтірілген. Параметрі бар есеп гиперболалық тендеулер жүйесі үшін параметрі бар бейлокал есептен және интегралдық қатынастан тұрады. Параметрі бар пара-пара есептің жуық шешімін табу алгоритмдері тұрғызылған және олардың жинақтылығы дәлелденген. Параметрі бар есептің жалғыз шешімінің бар болуының жеткілікті шарттары тағайындалған. Гиперболалық тектес интегралдық-дифференциалдық тендеулер жүйесі үшін бейлокал есептің жалғыз классикалық шешімінің бар болуының шарттары бастапқы берілімдер терминінде алынған. Қарастырылып отырған есепті зерттеу үшін бұрын дербес туындылы дифференциалдық тендеулер үшін есептер әулетіне келтіру әдісі пайдаланылған болатын. Зерттеліп отырған есептің жалғыз классикалық шешімінің бар болуының шарттары бастапқы берілімдер арқылы тұрғызылатын матрица терминінде табылған.

Тірек сөздер: бейлокал есеп, дербес туындылы интегралдық-дифференциалдық тендеулер жүйесі, параметр, алгоритм, жуық шешім, бірімәнді шешілімділік.

А.Т. Асанова, Х.А. Аширбаев, А.П. Сабалахова

О НЕЛОКАЛЬНОЙ ЗАДАЧЕ ДЛЯ СИСТЕМЫ ИНТЕГРО-ДИФФЕРЕНЦИАЛЬНЫХ УРАВНЕНИЙ В ЧАСТНЫХ ПРОИЗВОДНЫХ ГИПЕРБОЛИЧЕСКОГО ТИПА

Аннотация. Рассматривается нелокальная задача с данными на характеристиках для системы интегро-дифференциальных уравнений гиперболического типа второго порядка. Исследуются вопросы существования и единственности классического решения нелокальной задачи. Путем введения новой неизвестной функции вместо интегральной слагаемой исследуемая задача сведена к эквивалентной нелокальной задаче с интегральным условием. Задача с параметром состоит из нелокальной задачи для системы гиперболических уравнений с параметром и интегрального соотношения. Построены алгоритмы нахождения приближенного решения эквивалентной задачи с параметром и доказана их сходимость. Установлены достаточные условия существования единственного решения задачи с параметром. Получены условия существования единственного классического решения нелокальной задачи для системы интегро-дифференциальных уравнений гиперболического типа в терминах исходных данных. Ранее к исследованию рассматриваемой задачи был применен метод сведения к эквивалентному семейству задач для дифференциальных уравнений в частных производных. Были найдены достаточные условия существования единственного классического решения исследуемой задачи в терминах некоторой матрицы, составляемой по исходным данным.

Ключевые слова: нелокальная задача, система интегро-дифференциальных уравнений в частных производных, параметр, алгоритм, приближенное решение, однозначная разрешимость.

CONTENT

Technical sciences	
<i>Azamatov B.N., Ozhikenov K.A., Azamatova Zh. K.</i> ACS of the set of hydrocyclones with a variable geometry in the system of HAR TPP	5
Social Sciences	
<i>Kaldybay K.K., Pazylova K.A.</i> Socio-psychological analysis of the concept of aggression.....	14
Technical sciences	
<i>Aubakirova B.N., Beisenova R.R., Zhamangara A.K.</i> The effect of pharmaceutical ingredients to the growth of algae.....	21
<i>Saidullayeva N.S., Kabyrbekov K.A., Pazylova D.T., Ashirbaev Kh.A., Kalikulova A.O.</i> Designing the ambiguous tasks and tasks with missing data for performance of computer laboratory works.....	28
<i>Assanova A.T., Ashirbaev H.A., Sabalakhova A.P.</i> On the nonlocal problem for a system of the partial integro-differential equations of hyperbolic type.....	34
<i>Biyasheva Z.M., Zhumabai A.N., Shaizadinova A.M., Tleubergenova M.Zh., Sarzhanova S.D.</i> Mutagenic effects of alpha radiation in drosophil α test-system.....	43

**Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the work described has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the originality detection service Cross Check <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

www.nauka-nanrk.kz

ISSN 2518-1483 (Online), ISSN 2224-5227 (Print)

<http://www.reports-science.kz/index.php/ru/>

Редакторы *М. С. Ахметова, Д. С. Аленов, Т.А. Апендиев*
Верстка на компьютере *А.М. Кульгинбаевой*

Подписано в печать 15.08.2017.
Формат 60x881/8. Бумага офсетная. Печать – ризограф.
12,3 п.л. Тираж 2000. Заказ 4.

Национальная академия наук РК
050010, Алматы, ул. Шевченко, 28, т. 272-13-18, 272-13-19